

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 August 2005 (04.08.2005)

PCT

(10) International Publication Number
WO 2005/070900 A1

(51) International Patent Classification⁷: **C07D 239/42**,
409/14, 403/04, 405/04, A61K 31/505, 31/506, A61P
35/00, 37/00

(21) International Application Number:
PCT/EP2005/050206

(22) International Filing Date: 19 January 2005 (19.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04001310.4 22 January 2004 (22.01.2004) EP

(71) Applicant (for all designated States except US): **ALTANA PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BARSIG, Johannes** [DE/DE]; Bleichenweg 11, 78467 Konstanz (DE). **BAUDLER, Monika** [DE/DE]; Doebelestr. 16, 78462 Konstanz (DE). **BUNDSCHUH, Daniela** [DE/CH]; Im Rank 5, CH-8272 Ermatingen (CH). **GANTNER, Florian** [AT/CH]; Im Rank 5, CH-8272 Ermatingen (CH).

GRAEDLER, Ulrich [DE/DE]; Fichtestr. 15, 69126 Heidelberg (DE). **HEIT, Isabelle** [DE/CH]; Ruchwiesenstr. 44, CH-8404 Winterthur (CH). **MARTIN, Thomas** [DE/DE]; St.-Martins-Weg 13, 78462 Konstanz (DE). **SCHAEFER, Michaela** [DE/DE]; Friedhofstr. 1, 78359 Orsingen-Nenzingen (DE). **SCHLEMMINGER, Imre** [DE/DE]; Dorfbachweg 1, 78467 Konstanz (DE). **STADL-WIESER, Josef** [AT/DE]; Im Apfelgarten 3, 78465 Konstanz (DE). **ULRICH, Wolf-Ruediger** [DE/DE]; Alpenstr. 2, 78464 Konstanz (DE).



(74) Agents: **WILD, Robert** et al.; c/o ALTANA Pharma AG, 78467 Konstanz (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]

(54) Title: N-4-(6- (HETEO) ARYL-PYRIMIDIN-4-YLAMINOPHENYL) -BEZENESULFONAMIDES AS KINASE INHIBITORS



(57) Abstract: Compounds of a certain formula 1 in which R1 and R2 have the meanings indicated in the description are novel kinase inhibitors. Formula (1), in which R1 is phenyl, phenyl substituted by R3 and/or R4, naphthalenyl, naphthalenyl substituted by R5 and/or R6, aryl1, aryl1 substituted by R7 and/or R8, R9, R10 or R11, R2 is phenyl, phenyl substituted by R12 and/or R13, naphthalenyl, naphthalenyl substituted by R14 and/or R15, aryl2, aryl2 substituted by R16 and/or R17 or a radical selected from Formula (2).



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,*

NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- *of inventorship (Rule 4.17(iv)) for US only*

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

- 1 -

N-4-(6-(HETERO)ARYL-PYRAMIDIN-4-YLAMINOPHENYL)-BENZENESULFONAMIDES AS KINASE INHIBITORS**Field of application of the invention**

The invention relates to novel kinase inhibitors, which are used in the pharmaceutical industry for the production of pharmaceutical compositions.


Known technical background

In the International patent application WO02/12198 4-pyrimidineamine derivatives with neuroprotective properties are described.

Description of the invention

It has now been found that the kinase inhibitors, which are described in greater details below, have surprising and particularly advantageous properties.

The invention thus relates to compounds of formula 1




in which

- R1 is phenyl,
phenyl substituted by R3 and/or R4,
naphthalenyl,
naphthalenyl substituted by R5 and/or R6,
aryl1,
aryl1 substituted by R7 and/or R8,
R9,
R10 or
R11,
R2 is phenyl,

- 2 -

phenyl substituted by R12 and/or R13,
 naphthalenyl,
 naphthalenyl substituted by R14 and/or R15,
 aryl²,
 aryl² substituted by R16 and/or R17
 or a radical selected from



R3 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$ wherein

R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein

R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

n is an integer from 0 to 4,

m is an integer from 2 to 4,

R4 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,


R5 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl,

- 3 -

aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$,

- R6 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- Aryl1 furanyl, thiophenyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinoxalinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl or dibenzofuranyl,
- R7 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$,
- R8 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R9 is unsubstituted pyrrolyl, pyrazolyl, imidazolyl, indolyl, indazolyl, benzimidazolyl or benztriazolyl, or a radical selected from

- 4 -




wherein

R91 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$,

R92 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R10 is a radical selected from




wherein

R101 is 1-4C-alkyl, 2,2,2-trifluoroethyl or 3,3,3-trifluoropropyl,

- 5 -

R102 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R11 is a radical selected from



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R112)R113$ or $-CH_2CH(OH)CH_2N(R112)R113$, wherein

R112 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R113 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R112 and R113 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

- 6 -

Aryl² furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,

R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds with the proviso that the following compounds are excluded

4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(2-Benzoyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(4-Benzoyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl]-acetamide,
 N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
 N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
 N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide, and
 4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

- 7 -

Halogen within the meaning of the present invention is bromine, chlorine or fluorine.

1-4C-Alkyl is a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

1-4C-Alkoxy is a radical which, in addition to the oxygen atom, contains a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Alkoxy radicals having 1 to 4 carbon atoms which may be mentioned in this context are, for example, the butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy, ethoxy and methoxy radicals.

1-4C-Alkoxy which is completely or predominantly substituted by fluorine is, for example, the 2,2,3,3,3-pentafluoropropoxy, the perfluoroethoxy, the 1,2,2-trifluoroethoxy and in particular the 1,1,2,2-tetrafluoroethoxy, the 2,2,2-trifluoroethoxy, the trifluoromethoxy and the difluoromethoxy radical, of which the difluoromethoxy radical is preferred. "Predominantly" in this connection means that more than half of the hydrogen atoms of the 1-4C-alkoxy groups are replaced by fluorine atoms.

1-4C-Alkoxy carbonyl is a carbonyl group to which one of the abovementioned 1-4C-alkoxy radicals is bonded. Examples are the methoxycarbonyl $[\text{CH}_3\text{O}-\text{C}(\text{O})-]$ and the ethoxycarbonyl $[\text{CH}_3\text{CH}_2\text{O}-\text{C}(\text{O})-]$ radical.

1-4C-Alkyl carbonyl is a carbonyl group to which one of the abovementioned 1-4C-alkyl radicals is bonded. An example is the acetyl radical $[\text{CH}_3\text{CO}-]$.

Mono- or Di-1-4C-alkylamino radicals contain in addition to the nitrogen atom, one or two of the abovementioned 1-4C-alkyl radicals. Preferred are the di-1-4C-alkylamino radicals, especially the dimethylamino, the diethylamino and the diisopropylamino radical.

2-4C-Alkyl is a straight-chain or branched alkyl radical having 2 to 4 carbon atoms. Examples are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl and ethyl radicals.

1-4C-Alkoxy-2-4C-alkyl stands for one of the abovementioned 2-4C-alkyl radicals which is substituted by one of the abovementioned 1-4C-alkoxy radicals. Examples which may be mentioned are the 2-methoxyethyl and the 3-methoxypropyl radical.


Mono- or di-1-4C-alkylaminocarbonyl radicals are, for example, the methylaminocarbonyl, the dimethylaminocarbonyl and the diethylaminocarbonyl radicals.

- 8 -

An 1-4C-alkylcarbonylamino radical is, for example, the propionylamino [$\text{C}_3\text{H}_7\text{C}(\text{O})\text{NH}\cdot$] and the acetyl-amino radical [$\text{CH}_3\text{C}(\text{O})\text{NH}\cdot$].


$\cdot(\text{CH}_2)_n$, $\cdot(\text{CH}_2)_m$ or $\cdot(\text{CH}_2)_p$ stands for a straight-chain or branched alkylene radical having, n, m or p carbon atoms. Examples, which may be mentioned are methylene ($\cdot\text{CH}_2$), ethylene ($\cdot\text{CH}_2\text{CH}_2$), trimethylene ($\cdot\text{CH}_2\text{CH}_2\text{CH}_2$) or tetramethylene ($\cdot\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$).

In case R9 is a radical selected from




R91, R92 and the bond to the pyrimidinyl-ring can be attached to any carbon atom of the condensed ring systems with a free valence.

In case R10 is a radical selected from



R102 and the bond to the pyrimidinyl-ring can be attached to any carbon atom of the condensed ring systems with a free valence. Preferred are those cases, wherein R10 is a indol-5-yl or indazol-5-yl radical, and R102 is attached in 3-position or wherein R10 is a indol-3-yl or indazol-3-yl radical and R102 is attached in 5-position.

In case R11 is a radical selected from



R114 and the bond to the pyrimidinyl-ring can be attached to any carbon atom of the condensed ring systems with a free valence. Preferred are those cases, wherein R11 is a indol-5-yl or indazol-5-yl radical.

Suitable salts for compounds of the formula 1 - depending on substitution - are all acid addition salts or all salts with bases. Particular mention may be made of the pharmacologically tolerable inorganic and organic acids and bases customarily used in pharmacy. Those suitable are, on the one hand, water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulphuric acid, formic acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulphosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulphonic acid, methanesulphonic acid or 3-hydroxy-2-naphthoic acid, the acids being employed in salt preparation - depending on whether a mono- or polybasic acid is concerned and depending on which salt is desired - in an equimolar quantitative ratio or one differing therefrom.

On the other hand, salts with bases are - depending on substitution - also suitable. As examples of salts with bases are mentioned the lithium, sodium, potassium, calcium, aluminium, magnesium, titanium, ammonium, meglumine or guanidinium salts, here, too, the bases being employed in salt preparation in an equimolar quantitative ratio or one differing therefrom.


Pharmacologically intolerable salts, which can be obtained, for example, as process products during the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically tolerable salts by processes known to the person skilled in the art.

According to expert's knowledge the compounds of the invention as well as their salts may contain, e.g. when isolated in crystalline form, varying amounts of solvents. Included within the scope of the invention are therefore all solvates and in particular all hydrates of the compounds of formula 1 as well as all solvates and in particular all hydrates of the salts of the compounds of formula 1.

An embodiment (embodiment A) of the compounds of formula 1 are those in which

- R1 is phenyl,
phenyl substituted by R3 and/or R4,
naphthalenyl or
naphthalenyl substituted by R5 and/or R6,
- R2 is phenyl,
phenyl substituted by R12 and/or R13,
naphthalenyl,
naphthalenyl substituted by R14 and/or R15,
aryl2,
aryl2 substituted by R16 and/or R17
or a radical selected from

- 10 -



- R3 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy or benzyloxy,
- R4 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R5 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy or benzyloxy,
- R6 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-

- 11 -

dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,

- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds with the proviso that the following compounds are excluded

4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(2-Benzoyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(4-Benzoyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-(3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl)-acetamide,
 N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
 N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
 N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide, and
 N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide.

Compounds of formula 1 of embodiment A to be emphasized are those in which

- R1 is phenyl, 3-fluorophenyl, 4-fluorophenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-dimethoxy-phenyl, 3-acetylphenyl, 4-acetylphenyl, 3-cyanophenyl, 4-phenoxyphenyl or naphthalen-1-yl,
- R2 is phenyl, phenyl substituted by R12 and/or R13, thiophenyl, naphthalenyl or 5-(1-methyl-5-trifluoromethyl-1H-pyrazol-3-yl)-thiophene-2-yl,
- R12 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine or 1-4C-alkoxycarbonyl,
- R13 is halogen, 1-4C-alkyl or 1-4C-alkoxy,

and the salts of these compounds with the proviso that the following compounds are excluded

4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide, and
 N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide.

- 12 -

Preferred compounds of formula 1 of embodiment A are those in which

- R1 is phenyl, 3-fluorophenyl, 4-fluorophenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-acetylphenyl or naphthalen-1-yl,
 R2 is 2-fluorophenyl, 3-fluorophenyl, 3,4-difluorophenyl, 2,4-difluorophenyl, 2,6-difluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluorophenyl, 3-chloro-4-methylphenyl, 3-bromophenyl, 3-methylphenyl, 4-methylphenyl, 4-isopropylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 4-cyanophenyl or 5-(1-methyl-5-trifluoromethyl-1H-pyrazol-3-yl)-thiophene-2-yl,


and the salts of these compounds with the proviso that the following compounds are excluded

4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide and

4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

Another embodiment (embodiment B) of the compounds of formula 1 are those in which

- R1 is phenyl substituted by R3 and/or R4 or naphthalenyl substituted by R5 and/or R6,
 R2 is phenyl,
 phenyl substituted by R12 and/or R13,
 naphthalenyl,
 naphthalenyl substituted by R14 and/or R15,
 aryl2,
 aryl2 substituted by R16 and/or R17
 or a radical selected from



- R3 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-

- 13 -

alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein

R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein

R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

n is an integer from 0 to 4,

m is an integer from 2 to 4,

R4 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R5 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$,

R6 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,

- 14 -

- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- and the salts of these compounds.

Compounds of formula 1 of embodiment B to be emphasized are those in which

- R1 is phenyl substituted in para or meta position by R3 or phenyl substituted in para and meta position by R3 and R4,
- R2 is phenyl substituted by R12 and/or R13,
- R3 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein
- R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and
- R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein
- R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,
- n is an integer from 1 to 4,
- m is an integer from 2 to 4,
- R4 is fluorine, methyl or methoxy,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- and the salts of these compounds.

Further compounds of formula 1 of embodiment B to be emphasized are those in which

- R1 is phenyl substituted in para or meta position by R3,

- 15 -

- R2 is phenyl substituted by R12 and/or R13,
- R3 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein
- R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and
- R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein
- R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,
- n is an integer from 1 to 4,
- m is an integer from 2 to 4,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- and the salts of these compounds.

Compounds of formula 1 of embodiment B particularly to be emphasized are those in which

- R1 is phenyl substituted in para position by R3 or phenyl substituted in para position by R3 and in meta position by R4,
- R2 is phenyl substituted by R12 and/or R13,
- R3 is morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylmethyl, 4-methylpiperazin-1-ylethyl, 4-methylpiperazin-1-ylpropyl, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 1-methylpiperidin-4-ylmethoxy, 1-methylpiperidin-4-ylmethyl, 1-methylpiperidin-4-ylethyl, 1-methylpiperidin-4-ylpropyl, 2-(1-methylpiperidin-4-yl)ethoxy, 3-(1-methylpiperidin-4-yl)propoxy, pyrrolidin-1-ylpropyl, pyrrolidin-1-ylethyl, pyrrolidin-1-ylmethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy, piperidin-1-ylmethyl, piperidin-1-ylethyl, piperidin-1-ylpropyl, 2-(piperidin-1-yl)ethoxy, 3-(piperidin-1-yl)propoxy,
- R4 is fluorine, methyl or methoxy,
- R12 is fluorine, chlorine, cyano, methyl, isopropyl, trifluoromethyl or methoxy,

- 16 -

R13 is fluorine or chlorine,
and the salts of these compounds.

Further compounds of formula 1 of embodiment B particularly to be emphasized are those in which

- R1 is phenyl substituted in para or meta position by R3,
R2 is phenyl substituted by R12 and/or R13,
R3 is morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylmethyl, 4-methylpiperazin-1-ylethyl, 4-methylpiperazin-1-ylpropyl, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 1-methylpiperidin-4-ylmethoxy, 1-methylpiperidin-4-ylmethyl, 1-methylpiperidin-4-ylethyl, 1-methylpiperidin-4-ylpropyl, 2-(1-methylpiperidin-4-yl)ethoxy, 3-(1-methylpiperidin-4-yl)propoxy, pyrrolidin-1-ylpropyl, pyrrolidin-1-ylethyl, pyrrolidin-1-ylmethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy, piperidin-1-ylmethyl, piperidin-1-ylethyl, piperidin-1-ylpropyl, 2-(piperidin-1-yl)ethoxy, 3-(piperidin-1-yl)propoxy,
R12 is fluorine, chlorine, cyano, methyl, isopropyl, trifluoromethyl or methoxy,
R13 is fluorine or chlorine,
and the salts of these compounds.

Preferred compounds of formula 1 of embodiment B are those in which

- R1 is phenyl substituted in para position by R3 or phenyl substituted in para position by R3 and in meta position by R4,
R2 is 2-fluorophenyl, 4-fluorophenyl, 2,6-difluorophenyl, 2,4-difluorophenyl, 2-fluoro-4-methylphenyl, 2-fluoro-4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 4-methylphenyl, 4-methoxyphenyl or 3-chloro-4-fluorophenyl,
R3 is 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylethoxy, 4-methylpiperazin-1-ylpropoxy, morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 1-methylpiperidin-4-ylmethoxy, 2-(1-methylpiperidin-4-yl)ethoxy, 4-methylpiperazin-1-ylethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy or 3-pyrrolidin-1-ylpropyl,
R4 is fluorine,
and the salts of these compounds.

Further preferred compounds of formula 1 of embodiment B are those in which


- R1 is phenyl substituted in para or meta position by R3,
R2 is 2-fluorophenyl, 4-fluorophenyl, 2,6-difluorophenyl, 2,4-difluorophenyl, 4-methylphenyl, 4-methoxyphenyl or 3-chloro-4-fluorophenyl,
R3 is 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylethoxy, 4-methylpiperazin-1-ylpropoxy, morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl,

- 17 -

1-methylpiperidin-4-ylmethoxy, 2-(1-methylpiperidin-4-yl)ethoxy, 4-methylpiperazin-1-ylethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy or 3-pyrrolidin-1-ylpropyl, and the salts of these compounds.

A further embodiment (embodiment C) of the compounds of formula 1 are those in which

- R1 is aryl1,
 aryl1 substituted by R7 and/or R8,
 R9 or
 R10,
 R2 is phenyl,
 phenyl substituted by R12 and/or R13,
 naphthalenyl,
 naphthalenyl substituted by R14 and/or R15,
 aryl2,
 aryl2 substituted by R16 and/or R17
 or a radical selected from



- Aryl1 furanyl, thiophenyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl or dibenzofuranyl,
 R7 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyl-oxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-

- 18 -

piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R_{31})R_{32}$, $-CH_2CH(OH)CH_2N(R_{31})R_{32}$ or $-O-(CH_2)_m-N(R_{31})R_{32}$, wherein

R₃₁ is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R₃₂ is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein


R₃₁ and R₃₂ together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

n is an integer from 0 to 4,

m is an integer from 2 to 4,

R₈ is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R₉ is unsubstituted pyrrolyl, pyrazolyl, imidazolyl, indolyl, indazolyl, benzimidazolyl or benzotriazolyl, or a radical selected from




wherein

R₉₁ is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R_{31})R_{32}$, $-CH_2CH(OH)CH_2N(R_{31})R_{32}$ or $-O-(CH_2)_m-N(R_{31})R_{32}$,

R₉₂ is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

- 19 -

R10 is a radical selected from



wherein

R101 is 1-4C-alkyl, 2,2,2-trifluoroethyl or 3,3,3-trifluoropropyl,

R102 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-dioxinyl, quinoxalinyl, quinoxaliny, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,

R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

- 20 -

and the salts of these compounds with the proviso that the following compounds are excluded

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide, and
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

Compounds of formula 1 of embodiment C to be emphasized are those in which

- R1 is pyrid-3-yl, pyrid-4-yl, 2-methyl-2,3-dihydrobenzofuran-5-yl, benzo[1,3]dioxol-5-yl, 1-methyl-1H-pyrrol-3-yl, 4-methyl-thiophen-2-yl, 1-methyl-1H-pyrrol-2-yl, 1H-indol-5-yl, 1-methyl-1H-indol-3-yl, 1-methyl-1H-indol-5-yl, dibenzofuran-4-yl or 3,5-dimethyl-isoxazol-4-yl,
R2 is phenyl substituted by R12 and/or R13 or naphthalenyl,
R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds with the proviso that the following compounds are excluded

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide, and
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

Compounds of formula 1 of embodiment C particularly to be emphasized are those in which

- R1 is 2-methyl-2,3-dihydrobenzofuran-5-yl, benzo[1,3]dioxol-5-yl, 1-methyl-1H-pyrrol-3-yl, 4-methyl-thiophen-2-yl, 1-methyl-1H-pyrrol-2-yl, 1H-indol-5-yl, 1-methyl-1H-indol-3-yl or 1-methyl-1H-indol-5-yl,
R2 is phenyl substituted by R12 and/or R13 or naphthalenyl,
R12 is fluorine, chlorine, cyano, methyl, trifluoromethyl or methoxy,
R13 is fluorine, chlorine or methoxy,

and the salts of these compounds with the proviso that the compound

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide is excluded.

Preferred compounds of formula 1 of embodiment C are those in which

- 21 -

either

R1 is 2-methyl-2,3-dihydrobenzofuran-5-yl, 1-methyl-1H-pyrrol-3-yl or 4-methylthiophen-2-yl, and

R2 is 2,6-difluorophenyl,

or

R1 is 1H-indol-5-yl, and

R2 is 2,6-difluorophenyl, 2,4-difluorophenyl, 2-fluoro-4-methylphenyl, 2-fluorophenyl or 3-fluorophenyl,

or

R1 is 1-methyl-1H-indol-5-yl or 1-methyl-1H-indol-3-yl, and

R2 is 2,6-difluorophenyl, 2-fluorophenyl or 4-methoxyphenyl,

or

R1 is 1-methyl-1H-indol-5-yl, and

R2 is 2-fluoro-4-methylphenyl,

and the salts of these compounds.

Further preferred compounds of formula 1 of embodiment C are those in which

either

R1 is 2-methyl-2,3-dihydrobenzofuran-5-yl, 1-methyl-1H-pyrrol-3-yl or 4-methylthiophen-2-yl, and

R2 is 2,6-difluorophenyl,

or

R1 is 1H-indol-5-yl, and

R2 is 2,6-difluorophenyl, 2,4-difluorophenyl, 2-fluorophenyl or 3-fluorophenyl,

or

R1 is 1-methyl-1H-indol-5-yl or 1-methyl-1H-indol-3-yl, and

R2 is 2,6-difluorophenyl, 2-fluorophenyl or 4-methoxyphenyl,

and the salts of these compounds.

Still a further embodiment (embodiment D) of the compounds of formula 1 are those in which

R1 is R11,

R2 is phenyl,

phenyl substituted by R12 and/or R13,

naphthalenyl,


naphthalenyl substituted by R14 and/or R15,

aryl²,


aryl² substituted by R16 and/or R17

or a radical selected from

- 22 -



R11 is a radical selected from



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R112)R113$ or $-CH_2CH(OH)CH_2N(R112)R113$, wherein

R112 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R113 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R112 and R113 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,


- 23 -

- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,
- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds.

Compounds of formula 1 of embodiment D to be emphasized are those in which

- R1 is R11,
- R2 is phenyl substituted by R12 and/or R13,
- R11 is a radical selected from



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl,

- 24 -

1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R_{112})R_{113}$ or $-CH_2CH(OH)CH_2N(R_{112})R_{113}$, wherein

R_{112} is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R_{113} is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R_{112} and R_{113} together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R_{114} is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R_{12} is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R_{13} is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,


and the salts of these compounds.

Further compounds of formula 1 of embodiment D to be emphasized are those in which

R_1 is R_{11} ,

R_2 is phenyl substituted by R_{12} and/or R_{13} ,

R_{11} is a radical selected from



wherein

R_{111} is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R_{112})R_{113}$ or $-CH_2CH(OH)CH_2N(R_{112})R_{113}$, wherein

R_{112} is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R_{113} is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R_{112} and R_{113} together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

- 25 -

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,


and the salts of these compounds.

Compounds of formula 1 of embodiment D particularly to be emphasized are those in which

R1 is R11,

R2 is phenyl substituted by R12 and/or R13,

R11 is a radical selected from



wherein

R111 is pyrrolidin-1-ylethyl, pyrrolidin-1-ylpropyl, piperidin-1-ylethyl, piperidin-1-ylpropyl, azepan-1-ylethyl, azepan-1-ylpropyl, (4-methyl-piperazin-1-yl)ethyl, (4-methyl-piperazin-1-yl)propyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, (1-methyl-piperidin-4-yl)propyl, (1-methyl-piperidin-4-yl)ethyl, (1-methyl-piperidin-4-yl)methyl, dimethylaminoethyl, dimethylaminopropyl, dimethylaminobutyl, diethylaminoethyl, diethylaminopropyl or diethylaminobutyl,

R114 is hydrogen or fluorine,

R12 is fluorine, chlorine, cyano, methyl, isopropyl, trifluoromethyl or methoxy,

R13 is fluorine or chlorine,

and the salts of these compounds.


Preferred compounds of formula 1 of embodiment D are those in which

R1 is R11,

R2 is 2-fluorophenyl, 4-fluorophenyl, 2,6-difluorophenyl, 2,4-difluorophenyl, 4-methylphenyl, 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 2-fluoro-4-methylphenyl, or 3-chloro-4-fluorophenyl,

R11 represents the following radical

- 26 -



wherein

R111 is pyrrolidin-1-ylethyl, pyrrolidin-1-ylpropyl, (4-methyl-piperazin-1-yl)ethyl, (4-methyl-piperazin-1-yl)propyl, (1-methyl-piperidin-4-yl)ethyl, (1-methyl-piperidin-4-yl)methyl, dimethylaminoethyl, dimethylaminopropyl, diethylaminopropyl or diethylaminobutyl, and the salts of these compounds.

The compounds of formula 1 according to the invention can, for example, be prepared as described in reaction scheme 1.

The compounds of formula 1 according to the invention can be prepared, for example, starting from 3-oxo-propionic acid ester derivatives of formula 8, wherein R1 has the above-mentioned meanings and R stands, for example, for 1-4C-alkyl. In a first reaction step [step 1] the ester derivatives of formula 8 are reacted with thiourea to give the corresponding 2-mercapto-pyrimidin-4-ol derivatives of formula 7.

In the next two reaction steps [step 2, step 3] the mercapto group of the compounds of formula 7 is removed by treatment with Raney-Ni in ethanol (-> compounds of formula 6) and the hydroxyl group is exchanged by a chlorine atom using POCl_3 . The resulting 4-chloropyrimidine derivatives of formula 5 are then reacted with (4-aminophenyl)-carbamic acid tert butyl ester [step 4] to give [4-(pyrimidin-4-ylamino)-phenyl]-carbamic acid tert butyl ester derivatives of formula 4.

Alternatively compounds of formula 4 can be obtained in a two steps procedure. In a first reaction step [step 5] 4,6-dichloropyrimidine is reacted with (4-aminophenyl)-carbamic acid tert butyl ester to give [4-(6-chloro-pyrimidin-4-yl-amino)-phenyl]-carbamic acid tert-butyl ester. Conversion of this compound with boronic acids of formula 9a [step 6] or boronic acid esters of formula 9b (for example 4,4,5,5-tetramethyl-(R1)-[1,3,2]dioxaborolane or 5,5-dimethyl-(R1)-[1,3,2]dioxaborinane), wherein R1 has the above-mentioned meanings lead to compounds of formula 4.

In the next reaction step [step 7] the amino group of the compounds of formula 4 is deprotected by HCl gas in dioxane leading to compounds of formula 3. In the final reaction step [step 8] the N-pyrimidin-4-yl-benzene-1,4-diamine derivatives of formula 3 are reacted with sulfonyl chloride derivatives of formula 2, wherein R2 has the above-mentioned meanings to yield the compounds of formula 1.


In addition, compounds of formula 1 can be obtained starting from [4-(6-chloro-pyrimidin-4-yl-amino)-phenyl]-carbamic acid tert-butyl ester. The carbamic acid tert butyl ester can be deprotected by HCl gas

- 27 -

in dioxane [step 9] to give N-(6-chloropyrimidin-4-yl)-benzene-1,4-diamine after basic workup, which on its part is reacted with sulfonyl chloride derivatives of formula 2 [step 10], wherein R2 have the above-mentioned meanings. The resulting compounds of formula 10 are reacted in a final reaction step [step 11] with boronic acids of formula 9a or boronic acid esters of formula 9b (for example 4,4,5,5-tetramethyl-(R1)-[1,3,2]dioxaborolane or 5,5-dimethyl-(R1)-[1,3,2]dioxaborinane), wherein R1 has the above-mentioned meanings to yield the compounds of formula 1.

Further information for the preparation of boronic acid esters can be found, for example, in Murata et al., J Org Chem 2000, 65, 6458 or J Org Chem 1997, 62, 164.

Reaction Scheme 1:



Suitably, the conversions are carried out analogous to methods, which are familiar per se to the person skilled in the art, for example, in the manner which is described in the following examples.

It is known to the person skilled in the art that if there are a number of reactive centers on a starting or intermediate compound it may be necessary to block one or more reactive centers temporarily by protective groups in order to allow a reaction to proceed specifically at the desired reaction center. A detailed description for the use of a large number of proven protective groups is found, for example, in T.W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991.

The substances according to the invention are isolated and purified in a manner known per se, e.g. by distilling off the solvent in vacuo and recrystallizing the residue obtained from a suitable solvent or subjecting it to one of the customary purification methods, such as column chromatography on a suitable support material.


Salts are obtained by dissolving the free compound in a suitable solvent (for example a ketone like acetone, methylethylketone, or methylisobutylketone, an ether, like diethyl ether, tetrahydrofuran or dioxane, a chlorinated hydrocarbon, such as methylene chloride or chloroform, or a low molecular weight aliphatic alcohol, such as ethanol, isopropanol) which contains the desired acid, or to which the desired acid is then added. The salts are obtained by filtering, reprecipitating, precipitating with a non-solvent for the addition salt or by evaporating the solvent. Salts obtained can be converted by basification into the free compounds which, in turn, can be converted into salts. In this manner, pharmacologically non-tolerable salts can be converted into pharmacologically tolerable salts.

The following examples illustrate the invention in greater detail, without restricting it. As well, further compounds of formula 1, of which the preparation is explicitly not described, can be prepared in an analogous way or in a way which is known by a person skilled in the art using customary preparation methods.

In the examples, h stands for hour(s), calc for calculated, fnd for found and MS for mass spectroscopy. ¹H-NMR stands for proton nuclear magnetic resonance spectroscopy. The compounds, which are mentioned in the examples as well as their salts are preferred compounds of the invention.

Several of the examples are prepared as formic acid salts; in these cases the calculated total weight and the total molecular formula corresponds to the addition of the calculated total weight respectively total molecular formula of the indicated compound and the formic acid.


- 30 -

Examples**Final products****1. 2,6-Difluoro-N-[4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**

N-[6-(4-Fluoro-phenyl)-pyrimidin-4-yl]-benzene-1,4-diamine (compound A9, 280 mg) and 2,6-lutidine (150 μ L) is dissolved in a mixture of dioxane (10 ml), dimethylformamide (0.5 ml) and water (0.1 ml). To the well stirred mixture a solution of 2,6-Difluoro-benzenesulfonyl chloride (213 mg) in dioxane (5 ml) is added and stirring is continued for about 2 h at ambient temperature. Saturated aqueous NaCl solution (5 ml) is added, the organic layer is separated and filtered through a plug of neutral alumina. The filtrate is concentrated in vacuo. The residue is dissolved in ethyl acetate and again filtered through neutral alumina. The filtrate is concentrated in vacuo. After crystallization from ethyl acetate and tert-BuOMe 148 mg of pure product is obtained as off-white solid.

$^1\text{H-NMR}$ (DMSO- d_6) δ (ppm): 10.67 (s, 1H, -NH); 9.62 (s, 1H, -NH); 8.65 (s, 1H); 8.06 (dxd, $J_1 = 8.9$ Hz, $J_2 = 5.5$ Hz, 2H); 7.69 (m, 1H); 7.60 (d, $J = 8.9$ Hz, 2H); 7.31 (m, 2H + 2H); 7.14 (s, 1H); 7.12 (d, $J = 8.8$ Hz, 2H).

MS(ESI): calc: $\text{C}_{22}\text{H}_{15}\text{F}_3\text{N}_4\text{O}_2\text{S}$ (456.45) m/z : [MH $^+$] 457.1 [$\text{m/e} = 457.1$ (MH $^+$, 100%)].

2. 2,4-Difluoro-N-[4-[6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

A mixture of N-[4-(6-chloro-pyrimidin-4-ylamino)-phenyl]-2,4-difluoro-benzenesulfonamide (compound A11, 455 mg), 1-methyl-4-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenoxy]methyl-piperidine (compound A14, 331 mg), 2M aqueous Cs_2CO_3 (1.2 ml) and trans-dichlorobis(tricyclohexylphosphine)-

- 31 -

palladium (20 mg) in degassed dimethoxyethane and ethanol is heated in a microwave oven (Emrys Optimizer from PersonalChemistry) at 140°C in a sealed vial for 20 min. The reaction mixture is filtered through a plug of Extrelute®. The filtrate is concentrated in vacuo. The crude product is purified by preparative HPLC (gradient from 25% to 75% acetonitrile in water buffered with formic acid and ammonium formic acid salt). The combined product fractions are freeze dried to obtain the pure product as formic acid salt. The free base is precipitated from an aqueous solution by careful addition of aqueous ammonia. Crystallization from acetonitrile and water affords 372 mg of pure compound as colourless solid.

¹H-NMR(DMSO-d₆) δ (ppm): 9.49 (s, 1H, -NH); 8.60 (s, 1H); 7.95 (d, J = 8.9 Hz, 2H); 7.68 (m, 1H); 7.56 (d, J = 8.9 Hz, 2H); 7.24 (dxd J₁ = J₂ = 9.0 Hz, 2H); 7.09 (d, J = 5.9 Hz, 2H); 7.08 (s, 1H); 7.05 (d, J = 5.9 Hz, 2H); 3.90 (d, J = 5.8 Hz, 2H); 2.82 (~d, J ~ 11.1 Hz, 2H); 2.19 (s, 3H); 1.92 (~t, J ~ 11.5 Hz, 2H); 1.75 (~m, 2H + 1H); 1.43 (~m, 2H).

MS(ESI): calc: C₂₃ H₂₃ F₂ N₅ O₃ S (565.65) fnd:[MH⁺] 566.3 [m/e = 566.3 (MH⁺, 100%)]

The following compounds can be prepared analogously to the methods described above:

3. 5-(1-Methyl-5-trifluoromethyl-1H-pyrazol-3-yl)-thiophene-2-sulfonic acid [4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-amide

MS: calc: C₂₅ H₁₉ F₃ N₆ O₂ S₂ (556.59) fnd:[MH⁺] 557.2

4. 4-Cyano-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₃ H₁₇ N₅ O₂ S (427.49) fnd:[MH⁺] 428.2

5. 4-Isopropyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₅ H₂₄ N₄ O₂ S (444.56) fnd:[MH⁺] 445.2

6. 4-Methoxy-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₃ H₂₀ N₄ O₃ S (432.50) fnd:[MH⁺] 433.2

7. 2-Fluoro-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₂ H₁₇ F N₄ O₂ S (420.47) fnd:[MH⁺] 421.2

8. 2,6-Difluoro-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

- 32 -

MS: calc: C₂₂ H₁₆ F₂ N₄ O₂ S (438.46) fnd:[MH⁺] 439.1

9. 2,4-Difluoro-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₈ F₂ N₄ O₂ S (488.52) fnd:[MH⁺] 489.2

10. 3,4-Dichloro-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₈ Cl₂ N₄ O₂ S (521.43) fnd:[MH⁺] 521.1

11. 3-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₇ H₂₂ N₄ O₂ S (466.57) fnd:[MH⁺] 467.1

12. N-[4-(6-Naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-trifluoromethyl-benzenesulfonamide

MS: calc: C₂₇ H₁₉ F₃ N₄ O₂ S (520.54) fnd:[MH⁺] 521.1

13. 2-Fluoro-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₉ F N₄ O₂ S (470.53) fnd:[MH⁺] 471.2

14. 3-Bromo-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₉ Br N₄ O₂ S (531.43) fnd:[MH⁺] 533.0

15. 3-Chloro-4-methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzene-sulfonamide

MS: calc: C₂₇ H₂₁ Cl N₄ O₂ S (501.01) fnd:[MH⁺] 501.1

16. 2,6-Difluoro-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₈ F₂ N₄ O₂ S (488.52) fnd:[MH⁺] 489.1

17. N-[4-[6-(3-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-trifluoromethyl-benzene-sulfonamide

MS: calc: C₂₄ H₁₉ F₃ N₄ O₃ S (500.50) fnd:[MH⁺] 501.2

- 33 -

18. 2,4-Difluoro-N-[4-[6-(3-methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamideMS: calc: C₂₃ H₁₈ F₂ N₄ O₃ S (468.49) fnd:[MH⁺] 469.1**19. 4-Cyano-N-[4-[6-(3-methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₄ H₁₉ N₅ O₃ S (457.51) fnd:[MH⁺] 458.1**20. 2-Fluoro-N-[4-[6-(3-methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₃ H₁₉ F N₄ O₃ S (450.50) fnd:[MH⁺] 451.2**21. 2,4-Difluoro-N-[4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₂ H₁₅ F₃ N₄ O₂ S (456.45) fnd:[MH⁺] 457.1**22. 4-Cyano-cyclohexa-1,5-dienesulfonic acid [4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-amide**MS: calc: C₂₃ H₁₈ F N₅ O₂ S (447.49) fnd:[MH⁺] 446.1**23. 2-Fluoro-N-[4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₂ H₁₆ F₂ N₄ O₂ S (438.46) fnd:[MH⁺] 439.1**24. 2,4-Difluoro-N-[4-[6-(3-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₂ H₁₅ F₃ N₄ O₂ S (456.45) fnd:[MH⁺] 457.2**25. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-2,4-difluoro-benzenesulfonamide**MS: calc: C₂₄ H₁₈ F₂ N₄ O₃ S (480.50) fnd:[MH⁺] 481.3**26. 2,6-Difluoro-N-[4-[6-(3-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₂ H₁₅ F₃ N₄ O₂ S (456.45) fnd:[MH⁺] 457.2**27. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-2,6-difluoro-benzenesulfonamide**MS: calc: C₂₄ H₁₈ F₂ N₄ O₃ S (480.50) fnd:[MH⁺] 481.2

- 34 -

28. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-3-chloro-4-fluoro-benzenesulfonamide

MS: calc: C₂₄ H₁₈ Cl F N₄ O₃ S (496.95) fnd:[MH⁺] 497.2

29. 2-Fluoro-N-[4-[6-(4-methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₃ H₁₉ F N₄ O₃ S (450.50) fnd:[MH⁺] 451.2

30. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-3,4-difluoro-benzenesulfonamide

MS: calc: C₂₄ H₁₈ F₂ N₄ O₃ S (480.50) fnd:[MH⁺] 481.2

31. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-3-fluoro-benzenesulfonamide

MS: calc: C₂₄ H₁₉ F N₄ O₃ S (462.51) fnd:[MH⁺] 463.2

32. 3-Fluoro-N-[4-[6-(4-methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₃ H₁₉ F N₄ O₃ S (450.50) fnd:[MH⁺] 451.2

33. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-3-methyl-benzenesulfonamide

MS: calc: C₂₅ H₂₂ N₄ O₃ S (458.54) fnd:[MH⁺] 459.2

34. N-[4-[6-(3-Fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide

MS: calc: C₂₃ H₁₉ F N₄ O₂ S (434.50) fnd:[MH⁺] 435.2

35. N-[4-[6-(4-Acetyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide

MS: calc: C₂₅ H₂₂ N₄ O₃ S (458.54) fnd:[MH⁺] 459.3

36. 2,6-Difluoro-N-[4-[6-(4-methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₃ H₁₈ F₂ N₄ O₃ S (468.49) fnd:[MH⁺] 469.2

37. 2,6-Difluoro-N-[4-[6-[4-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₈ H₂₇ F₂ N₅ O₄ S (567.62) fnd:[MH⁺] 568.3

38. 2,6-Difluoro-N-(4-{6-[4-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₉ H₂₉ F₂ N₅ O₄ S (567.62) fnd:[MH⁺] 582.3

39. 2,6-Difluoro-N-(4-{6-[3-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₉ H₂₉ F₂ N₅ O₄ S (581.65) fnd:[MH⁺] 582.3

40. 2-Fluoro-N-(4-{6-[4-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₈ H₂₈ F N₅ O₄ S (549.63) fnd:[MH⁺] 550.4

41. 2-Fluoro-N-(4-{6-[4-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₉ H₃₀ F N₅ O₄ S (563.66) fnd:[MH⁺] 564.4

42. 2-Fluoro-N-(4-{6-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₈ H₂₈ F N₅ O₄ S (549.93) fnd:[MH⁺] 550.4

43. 2-Fluoro-N-(4-{6-[3-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₉ H₃₀ F N₅ O₄ S (563.66) fnd:[MH⁺] 564.4

44. 4-Methoxy-N-(4-{6-[4-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₉ H₃₁ N₅ O₅ S (561.66) fnd:[MH⁺] 562.4

- 36 -

45. 4-Methoxy-N-[4-{6-[4-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₃ N₅ O₅ S (575.69) fnd:[MH⁺] 576.4

46. 4-Methoxy-N-[4-{6-[3-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₃ N₅ O₅ S (575.69) fnd:[MH⁺] 576.4

47. 2,6-Difluoro-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₉ H₃₀ F₂ N₆ O₃ S (580.66) fnd:[MH⁺] 581.4

48. 2,6-Difluoro-N-[4-(6-[3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₉ H₃₀ F₂ N₆ O₃ S (580.66) fnd:[MH⁺] 581.4

49. 2,6-Difluoro-N-[4-(6-[3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₂ F₂ N₆ O₃ S (594.69) fnd:[MH⁺] 595.4

50. 2-Fluoro-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₉ H₃₁ F N₆ O₃ S (562.67) fnd:[MH⁺] 563.3

51. 2-Fluoro-N-[4-(6-[3-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₃ F N₆ O₃ S (576.70) fnd:[MH⁺] 577.4

52. 4-Methoxy-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₄ N₆ O₄ S (574.71) fnd:[MH⁺] 575.4

- 37 -

53. 2,4-Difluoro-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₉ H₃₀ F₂ N₆ O₃ S (580.66) fnd:[MH⁺] 581.4

54. 4-Methyl-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₄ N₆ O₃ S (558.71) fnd:[MH⁺] 595.4

55. 4-Methyl-N-[4-(6-[4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₁ H₃₆ N₆ O₃ S (572.73) fnd:[MH⁺] 573.4

56. 3-Chloro-4-fluoro-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₉ H₃₀ Cl F N₆ O₃ S (597.12) fnd:[MH⁺] 597.4

57. 3-Chloro-4-fluoro-N-[4-(6-[4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₀ H₃₂ Cl F N₆ O₃ S (611.14) fnd:[MH⁺] 611.4

58. 2,4-Difluoro-N-(4-[6-[4-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide

MS: calc: C₂₉ H₂₉ F₂ N₆ O₄ S (581.65) fnd:[M+NH₄⁺] 598.5

59. 2,4-Difluoro-N-(4-[6-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide

MS: calc: C₂₈ H₂₇ F₂ N₅ O₄ S (567.62) fnd:[M+NH₄⁺] 584.4

60. 4-Methyl-N-(4-[6-[4-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide

MS: calc: C₃₀ H₃₃ N₅ O₄ S (559.69) fnd:[MH⁺] 560.4

- 38 -

61. 4-Methyl-N-(4-{6-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₃H₃₁N₅O₄S (545.67) fnd:[MH⁺] 546.4

62. 3-Chloro-4-fluoro-N-(4-{6-[4-(2-morpholin-4-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₈H₂₇ClF N₅O₄S (584.07) fnd:[MH⁺] 584.5

63. 3-Chloro-4-fluoro-N-(4-{6-[4-(3-morpholin-4-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide

MS: calc: C₂₉H₂₉ClF N₅O₄S (598.10) fnd:[MH⁺] 598.5

64. 2-Fluoro-N-(4-{6-(4-morpholin-4-ylmethyl-phenyl)-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₂₈H₂₈FN₅O₅S (565.63) fnd:[MH⁺] 520.2

65. 2-Fluoro-N-(4-{6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₃FN₆O₅S (608.70) fnd:[MH⁺] 563.2

66. 2-Fluoro-N-(4-{6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₄FN₅O₅S (607.71) fnd:[MH⁺] 562.2

67. 2-Fluoro-N-(4-{6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂FN₅O₅S (593.68) fnd:[MH⁺] 548.1

68. 2-Fluoro-N-(4-{6-[4-(2-morpholin-4-yl-ethyl)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₂₉H₃₀FN₅O₅S (579.66) fnd:[MH⁺] 534.3

- 39 -

69. 2-Fluoro-N-(4-{6-[4-(3-morpholin-4-yl-propyl)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂FN₅O₅S (593.68) fnd:[MH⁺] 548.2

70. 2-Fluoro-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₄FN₅O₅S (607.71) fnd:[MH⁺] 562.3

71. 2,6-Difluoro-N-(4-{6-[4-(3-pyrrolidin-1-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁F₂N₅O₅S (611.67) fnd:[MH⁺] 566.1

72. 2,6-Difluoro-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₃F₂N₅O₅S (625.70) fnd:[MH⁺] 580.2

73. 2,6-Difluoro-N-(4-{6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁F₂N₅O₅S (611.67) fnd:[MH⁺] 566.1

74. 2,6-Difluoro-N-(4-{6-[4-(3-morpholin-4-yl-propyl)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁F₂N₅O₅S (611.67) fnd:[MH⁺] 566.3

75. 2,6-Difluoro-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₃F₂N₅O₅S (625.70) fnd:[MH⁺] 580.2

76. 4-Methoxy-N-(4-{6-[4-(3-pyrrolidin-1-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₅N₅O₅S (605.72) fnd:[MH⁺] 560.3

- 40 -

77. 4-Methoxy-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₇N₅O₆S (619.75) fnd:[MH⁺] 574.3

78. 4-Methyl-N-(4-[6-[4-(3-pyrrolidin-1-yl-propoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₅N₅O₅S (589.72) fnd:[MH⁺] 544.3

79. 4-Methyl-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₇N₅O₅S (603.75) fnd:[MH⁺] 558.2

80. 4-Methyl-N-(4-[6-[4-(2-morpholin-4-yl-ethyl)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₃N₅O₅S (575.69) fnd:[MH⁺] 530.3

81. 4-Methyl-N-(4-[6-[4-(3-morpholin-4-yl-propyl)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₅N₅O₅S (589.72) fnd:[MH⁺] 544.3

82. 4-Methyl-N-(4-[6-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₃N₅O₅S (575.69) fnd:[MH⁺] 530.3

83. 4-Methyl-N-(4-[6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₅N₅O₅S (589.72) fnd:[MH⁺] 544.3

84. 4-Methyl-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₇N₅O₅S (603.75) fnd:[MH⁺] 558.3

- 41 -

85. 4-Methyl-N-(4-[6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₅N₆O₄S (612.76) fnd:[MH⁺] 567.3

86. 4-Methyl-N-[4-(6-[1-[2-(4-methyl-piperazin-1-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₇N₇O₄S (627.77) fnd:[MH⁺] 582.4

87. 2,4-Difluoro-N-(4-[6-[4-(3-pyrrolidin-1-yl-propoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁F₂N₅O₅S (611.67) fnd:[MH⁺] 566.2

88. 2,4-Difluoro-N-[4-[6-(4-morpholin-4-ylmethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₂₈H₂₇F₂N₅O₅S (583.62) fnd:[MH⁺] 538.1

89. 2,4-Difluoro-N-[4-(6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂F₂N₆O₅S (626.69) fnd:[MH⁺] 581.2

90. 2,4-Difluoro-N-[4-(6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂F₂N₆O₅S (625.70) fnd:[MH⁺] 580.2

91. 2,4-Difluoro-N-(4-[6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁F₂N₅O₅S (611.67) fnd:[MH⁺] 566.2

92. 2,4-Difluoro-N-(4-[6-[4-(2-morpholin-4-yl-ethyl)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₂₈H₂₉F₂N₅O₅S (597.65) fnd:[MH⁺] 552.2

- 42 -

93. 2,4-Difluoro-N-(4-[6-[4-(3-morpholin-4-yl-propyl)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁F₂N₅O₅S (611.67) fnd:[MH⁺] 566.3

94. 2,4-Difluoro-N-(4-[6-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₂₉H₂₉F₂N₅O₅S (597.65) fnd:[MH⁺] 552.2

95. 3-Chloro-4-fluoro-N-(4-[6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂ClFN₆O₅S (643.14) fnd:[MH⁺] 597.3

96. 3-Chloro-4-fluoro-N-(4-[6-[4-(3-morpholin-4-yl-propyl)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₁ClFN₅O₅S (628.13) fnd:[MH⁺] 582.3

97. 3-Chloro-4-fluoro-N-(4-[6-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₂₉H₂₉ClFN₅O₅S (614.10) fnd:[MH⁺] 568.3

98. 2-Fluoro-N-(4-[6-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₂₉H₃₀FN₅O₅S (579.66) fnd:[MH⁺] 534.2

99. 2-Fluoro-N-(4-[6-[4-(3-pyrrolidin-1-yl-propoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂FN₅O₅S (593.68) fnd:[MH⁺] 548.2

100. 2,6-Difluoro-N-(4-[6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂F₂N₆O₅S (626.69) fnd:[MH⁺] 581.2

- 43 -

101. 2,6-Difluoro-N-[4-{6-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{29}H_{29}F_2N_5O_5S$ (597.65) fnd: $[MH^+]$ 552.2

102. 4-Methoxy-N-[4-{6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{31}H_{36}N_6O_6S$ (620.73) fnd: $[MH^+]$ 575.3

103. 4-Methoxy-N-[4-{6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{31}H_{35}N_5O_6S$ (605.72) fnd: $[MH^+]$ 560.2

104. 4-Methoxy-N-[4-{6-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{30}H_{33}N_5O_6S$ (591.69) fnd: $[MH^+]$ 546.3

105. 4-Methyl-N-[4-{6-[4-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{31}H_{36}N_6O_5S$ (604.73) fnd: $[MH^+]$ 559.3

106. 4-Methyl-N-[4-{6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{31}H_{35}N_5O_5S$ (589.72) fnd: $[MH^+]$ 544.2

107. 4-Fluoro-N-[4-{6-[4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{31}H_{34}FN_5O_5S$ (607.71) fnd: $[MH^+]$ 562.2

108. 4-Fluoro-N-[4-{6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{30}H_{32}FN_5O_5S$ (593.68) fnd: $[MH^+]$ 548.2

- 44 -

109. 4-Fluoro-N-[4-{6-[4-(3-pyrrolidin-1-yl-propoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₀H₃₂FN₅O₅S (593.68) fnd:[MH⁺] 548.3

110. 2-Fluoro-N-[4-{6-[4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₅F N₆O₅S (622.72) fnd:[MH⁺] 577.3

111. 2,6-Difluoro-N-[4-{6-[4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₄F₂N₆O₅S (640.71) fnd:[MH⁺] 595.3

112. 2,4-Difluoro-N-[4-{6-[4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₄F₂N₆O₅S (640.71) fnd:[MH⁺] 595.3

113. 4-Fluoro-N-[4-{6-[4-[3-(4-methyl-piperazin-1-yl)-propoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₅F N₆O₅S (622.72) fnd:[MH⁺] 577.3

114. 2,4-Difluoro-N-[4-{6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₂₉H₂₉F₂N₅O₃S (565.65) fnd:[MH⁺] 566.2

115. 2,6-Difluoro-N-[4-{6-[2-methyl-2,3-dihydro-benzofuran-5-yl]-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₅H₂₀F₂N₄O₃S (494.52) fnd:[MH⁺] 495.2

116. 2,6-Difluoro-N-[4-{6-(1-methyl-1H-pyrrol-3-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₁H₁₇F₂N₅O₂S (441.46) fnd:[MH⁺] 442.0

- 45 -

117. 2,6-Difluoro-N-[4-[6-(4-methyl-thiophen-2-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₁ H₁₆ F₂ N₄ O₂ S₂ (458.51) fnd:[MH⁺] 459.0

118. 2,4-Difluoro-N-[4-[6-(1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₄ H₁₇ F₂ N₅ O₂ S (477.50) fnd:[MH⁺] 478.2

119. 2,6-Difluoro-N-[4-[6-(1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₄ H₁₇ F₂ N₅ O₂ S (477.50) fnd:[MH⁺] 478.2

120. 2-Fluoro-N-[4-[6-(1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₄ H₁₈ F N₅ O₂ S (459.51) fnd:[MH⁺] 460.2

121. 3-Fluoro-N-[4-[6-(1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₄ H₁₈ F N₅ O₂ S (459.51) fnd:[MH⁺] 460.2

122. 2,6-Difluoro-N-[4-[6-(1-methyl-1H-indol-3-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₉ F₂ N₅ O₂ S (491.52) fnd:[MH⁺] 492.3

123. 2,6-Difluoro-N-[4-[6-(1-methyl-1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₁₉ F₂ N₅ O₂ S (491.52) fnd:[MH⁺] 492.3

124. 2-Fluoro-N-[4-[6-(1-methyl-1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₂₀ F N₅ O₂ S (473.53) fnd:[MH⁺] 474.3

125. 4-Methoxy-N-[4-[6-(1-methyl-1H-indol-3-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide

MS: calc: C₂₆ H₂₃ N₅ O₃ S (485.57) fnd:[MH⁺] 486.3

- 46 -

126. 4-Methoxy-N-[4-[6-(1-methyl-1H-indol-5-yl)-pyrimidin-4-ylamino]-phenyl]-benzene-sulfonamideMS: calc: C₂₆ H₂₃ N₅ O₃ S (485.57) fnd:[MH⁺] 486.3**127. 2-Fluoro-N-[4-[6-(1-methyl-1H-indol-3-yl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**MS: calc: C₂₅ H₂₀ F N₅ O₂ S (473.53) fnd:[MH⁺] 474.3**128. N-(4-[6-[1-(3-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-2,6-difluoro-benzenesulfonamide**MS: calc: C₂₉ H₂₈ F₂ N₆ O₂ S (562.65) fnd:[MH⁺] 563.3**129. N-(4-[6-[1-(3-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-2-fluoro-benzenesulfonamide**MS: calc: C₂₉ H₂₉ F N₆ O₂ S (544.66) fnd:[MH⁺] 545.3**130. N-(4-[6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-2-fluoro-benzenesulfonamide**MS: calc: C₂₈ H₂₇ F N₆ O₂ S (530.63) fnd:[MH⁺] 531.3**131. N-(4-[6-[1-(3-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-4-methoxy-benzenesulfonamide**MS: calc: C₃₀ H₃₂ N₆ O₃ S (556.69) fnd:[MH⁺] 557.4**132. N-(4-[6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-4-methoxy-benzenesulfonamide**MS: calc: C₂₉ H₃₀ N₆ O₃ S (542.66) fnd:[MH⁺] 543.4**133. N-(4-[6-[1-(3-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-2,4-difluoro-benzenesulfonamide**MS: calc: C₂₉ H₂₈ F₂ N₆ O₂ S (562.65) fnd:[M+NH₄⁺] 579.3

- 47 -

134. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,4-difluoro-benzenesulfonamide

MS: calc: C₂₈ H₂₆ F₂ N₆ O₂ S (548.62) fnd: [M+NH₄⁺] 565.4

135. N-(4-{6-[1-(3-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide

MS: calc: C₃₀ H₃₂ N₆ O₂ S (540.69) fnd: [MH⁺] 541.4

136. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide

MS: calc: C₂₉ H₃₀ N₆ O₂ S (526.67) fnd: [MH⁺] 527.4

137. 3-Chloro-N-(4-{6-[1-(3-dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide

MS: calc: C₂₉ H₂₈ Cl F N₆ O₂ S (579.10) fnd: [MH⁺] 579.4

138. 3-Chloro-N-(4-{6-[1-(2-dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide

MS: calc: C₂₈ H₂₆ Cl F N₆ O₂ S (565.07) fnd: [MH⁺] 565.3

139. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide

MS: calc: C₂₈ H₂₆ F₂ N₆ O₂ S (548.62) fnd: [MH⁺] 549.3

140. 2-Fluoro-N-(4-{6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₁FN₆O₄S (602.69) fnd: [MH⁺] 557.3

141. 2-Fluoro-N-[4-{6-[1-[2-(4-methyl-piperazin-1-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₄FN₇O₄S (631.73) fnd: [MH⁺] 586.3

- 48 -

142. 2,6-Difluoro-N-(4-{6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₀F₂N₆O₄S (620.68) fnd:[MH⁺] 575.2

143. 2,6-Difluoro-N-(4-{6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₂F₂N₆O₄S (634.71) fnd:[MH⁺] 589.3

144. 2,6-Difluoro-N-[4-{6-[1-[2-(4-methyl-piperazin-1-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₃F₂N₇O₄S (649.73) fnd:[MH⁺] 604.3

145. 4-Methoxy-N-(4-{6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₆N₆O₅S (628.76) fnd:[MH⁺] 583.3

146. 4-Methyl-N-(4-{6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₆N₆O₄S (612.76) fnd:[MH⁺] 567.3

147. 4-Methyl-N-[4-{6-[1-[2-(4-methyl-piperazin-1-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₇N₇O₄S (627.77) fnd:[MH⁺] 582.4

148. 2,4-Difluoro-N-(4-{6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₀F₂N₆O₄S (620.68) fnd:[MH⁺] 575.3

149. 2,4-Difluoro-N-(4-{6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₂F₂N₆O₄S (634.71) fnd:[MH⁺] 589.3

- 49 -

150. 2,4-Difluoro-N-[4-(6-[1-[2-(4-methyl-piperazin-1-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{32}H_{33}F_2N_7O_4S$ (649.73) fnd: $[MH^+]$ 604.3

151. 4-Fluoro-N-(4-[6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{32}H_{33}FN_6O_4S$ (616.72) fnd: $[MH^+]$ 571.3

152. 2-Fluoro-N-(4-[6-[1-(3-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{32}H_{33}FN_6O_4S$ (616.72) fnd: $[MH^+]$ 571.3

153. 2-Fluoro-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{33}H_{36}FN_7O_4S$ (645.76) fnd: $[MH^+]$ 600.3

154. 2,6-Difluoro-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{33}H_{35}F_2N_7O_4S$ (663.75) fnd: $[MH^+]$ 618.3

155. 4-Methoxy-N-(4-[6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{32}H_{34}N_6O_5S$ (614.73) fnd: $[MH^+]$ 569.3

156. 4-Methoxy-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{34}H_{39}N_7O_5S$ (657.80) fnd: $[MH^+]$ 612.3

157. 4-Methoxy-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: $C_{33}H_{37}N_7O_5S$ (643.77) fnd: $[MH^+]$ 598.3

- 50 -

158. 4-Methyl-N-[4-(6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₂H₃₄N₆O₄S (598.73) fnd:[MH⁺] 553.3

159. 4-Methyl-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₄H₃₉N₇O₄S (641.80) fnd:[MH⁺] 596.3

160. 2,4-Difluoro-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₅F₂N₇O₄S (663.75) fnd:[MH⁺] 618.3

161. 3-Chloro-4-fluoro-N-[4-(6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁H₃₀ClFN₆O₄S (637.14) fnd:[MH⁺] 591.2

162. 3-Chloro-4-fluoro-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃H₃₅ClFN₇O₄S (680.21) fnd:[MH⁺] 634.3

163. 2-Fluoro-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-3-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₆ F N₇ O₄ S (645.76) fnd:[MH⁺] 600.3

164. 2,4-Difluoro-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-3-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₅ F₂ N₇ O₄ S (663.75) fnd:[MH⁺] 618.4

165. 4-Methoxy-N-[4-(6-[1-[3-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-3-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₄ H₃₉ N₇ O₅ S (657.80) fnd:[MH⁺] 612.3

- 51 -

166. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₄ F₂ N₆ O₄ S (636.73) fnd:[MH⁺] 591.3

167. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F N₆ O₄ S (618.74) fnd:[MH⁺] 573.3

168. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,4-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₄ F₂ N₆ O₄ S (636.73) fnd:[MH⁺] 591.3

169. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F N₆ O₄ S (618.74) fnd:[MH⁺] 573.3

170. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₈ N₆ O₅ S (630.77) fnd:[MH⁺] 585.3

171. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₈ N₆ O₄ S (614.77) fnd:[MH⁺] 569.3

172. N-(4-{6-[1-(3-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,4-difluoro-benzenesulfonamide

MS: calc: C₂₉ H₂₈ F₂ N₆ O₂ S (562.65) fnd:[MH⁺] 563.3

173. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-3-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₄ F₂ N₆ O₄ S (636.73) fnd:[MH⁺] 591.4

- 52 -

174. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-3-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F N₆ O₄ S (618.74) fnd:[MH⁺] 573.4

175. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-3-yl]-pyrimidin-4-ylamino}-phenyl)-2,4-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₄ F₂ N₆ O₄ S (636.73) fnd:[MH⁺] 591.3

176. N-(4-{6-[1-(3-Diethylamino-propyl)-1H-indol-3-yl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₈ N₆ O₅ S (630.77) fnd:[MH⁺] 585.4

177. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F N₆ O₄ S (604.71) fnd:[MH⁺] 559.3

178. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₆ N₆ O₄ S (600.75) fnd:[MH⁺] 555.3

179. 3-Chloro-N-(4-{6-[1-(2-diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₂ Cl F N₆ O₄ S (639.15) fnd:[MH⁺] 593.2

180. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,4-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₂ F₂ N₆ O₄ S (622.7) fnd:[MH⁺] 577.3

181. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₂ F₂ N₆ O₄ S (622.7) fnd:[MH⁺] 577.3

- 53 -

182. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₆ N₆ O₅ S (616.74) fnd:[MH⁺] 571.3

183. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F N₆ O₄ S (604.71) fnd:[MH⁺] 559.3

184. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-3-fluoro-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F N₆ O₅ S (634.74) fnd:[MH⁺] 589.3

185. N-(4-{6-[1-(2-Diethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F N₆ O₄ S (618.74) fnd:[MH⁺] 573.3

186. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₇ F N₆ O₄ S (632.76) fnd:[MH⁺] 587.4

187. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₄ H₄₀ N₆ O₄ S (628.8) fnd:[MH⁺] 583.4

188. 3-Chloro-N-(4-{6-[1-(4-diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₅ Cl F N₆ O₄ S (667.21) fnd:[MH⁺] 621.3

189. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,4-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₆ F₂ N₆ O₄ S (650.75) fnd:[MH⁺] 605.3

- 54 -

190. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₆ F₂ N₆ O₄ S (650.75) fnd:[MH⁺] 605.3

191. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₄ H₄₀ N₆ O₅ S (644.8) fnd:[MH⁺] 599.4

192. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₇ F N₆ O₄ S (632.76) fnd:[MH⁺] 587.4

193. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-3-fluoro-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₄ H₃₉ F N₆ O₅ S (662.79) fnd:[MH⁺] 617.3

194. N-(4-{6-[1-(4-Diethylamino-butyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₄ H₃₉ F N₆ O₄ S (646.79) fnd:[MH⁺] 601.3

195. 2-Fluoro-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₃ F N₆ O₄ S (616.72) fnd:[MH⁺] 571.3

196. 4-Methyl-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₆ N₆ O₄ S (612.76) fnd:[MH⁺] 567.4

197. 3-Chloro-4-fluoro-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₂ Cl F N₆ O₄ S (651.17) fnd:[MH⁺] 605.3

- 55 -

198. 2,4-Difluoro-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₂ F₂ N₆ O₄ S (634.71) fnd:[MH⁺] 589.3

199. 2,6-Difluoro-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₂ F₂ N₆ O₄ S (634.71) fnd:[MH⁺] 507.6

200. 4-Methoxy-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₆ N₆ O₅ S (628.76) fnd:[MH⁺] 583.4

201. 4-Fluoro-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₃ F N₆ O₄ S (616.72) fnd:[MH⁺] 571.3

202. 3-Fluoro-4-methoxy-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₅ F N₆ O₅ S (646.75) fnd:[MH⁺] 601.3

203. 2-Fluoro-4-methyl-N-(4-{6-[1-(1-methyl-piperidin-4-ylmethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₅ F N₆ O₄ S (630.75) fnd:[MH⁺] 585.3

204. 2-Fluoro-N-[4-(6-[1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₅ F N₆ O₄ S (630.75) fnd:[MH⁺] 585.3

205. 4-Methyl-N-[4-(6-[1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₃ H₃₆ N₆ O₂ S (580.76) fnd:[MH⁺] 581.3

- 56 -

206. 3-Chloro-4-fluoro-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₄ Cl F N₆ O₄ S (665.19) fnd:[MH⁺] 619.3

207. 2,4-Difluoro-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₄ F₂ N₆ O₄ S (648.74) fnd:[MH⁺] 603.3

208. 2,6-Difluoro-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₄ F₂ N₆ O₄ S (648.74) fnd:[MH⁺] 603.3

209. 4-Methoxy-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₄ H₃₈ N₆ O₅ S (642.78) fnd:[MH⁺] 597.3

210. 4-Fluoro-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₂ H₃₃ F N₆ O₂ S (584.72) fnd:[MH⁺] 585.3

211. 3-Fluoro-4-methoxy-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₃ H₃₅ F N₆ O₃ S (614.75) fnd:[MH⁺] 615.3

212. 2-Fluoro-4-methyl-N-[4-(6-{1-[2-(1-methyl-piperidin-4-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide

MS: calc: C₃₃ H₃₅ F N₆ O₂ S (598.75) fnd:[MH⁺] 599.3

213. 2,4-Difluoro-N-(4-{6-[3-fluoro-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino)-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₀ F₃ N₅ O₅ S (629.66) fnd:[MH⁺] 584.1

- 57 -

214. 3-Fluoro-N-[4-(6-[3-fluoro-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino)-phenyl]-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F₂ N₅ O₆ S (641.70) fnd:[MH⁺] 596.2

215. 2-Fluoro-N-[4-(6-[3-fluoro-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F₂ N₅ O₅ S (625.70) fnd:[MH⁺] 580.2

216. 2-Fluoro-N-[4-(6-[3-fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F₂ N₅ O₅ S (625.70) fnd:[MH⁺] 580.2

217. 2,4-Difluoro-N-[4-(6-[3-fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₂ F₃ N₅ O₅ S (643.69) fnd:[MH⁺] 598.2

218. 2,6-Difluoro-N-[4-(6-[3-fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₂ F₃ N₅ O₅ S (643.69) fnd:[MH⁺] 598.2

219. N-[4-(6-[3-Fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F N₅ O₆ S (637.74) fnd:[MH⁺] 592.2

220. 3-Fluoro-N-[4-(6-[3-fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F₂ N₅ O₆ S (655.73) fnd:[MH⁺] 610.2

221. 4-Fluoro-N-[4-(6-[3-fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

- 58 -

MS: calc: C₃₁ H₃₃ F₂ N₅ O₅ S (625.70) fnd:[MH⁺] 580.3

222. 2-Fluoro-N-[4-(6-[3-fluoro-4-[2-(1-methyl-piperidin-4-yl)-ethoxy]-phenyl]-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₅ F₂ N₅ O₅ S (639.73) fnd:[MH⁺] 594.2

223. 2-Fluoro-4-methyl-N-(4-[6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₄ F N₅ O₅ S (607.71) fnd:[MH⁺] 562.2

224. 3-Fluoro-4-methoxy-N-(4-[6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₄ F N₅ O₆ S (623.71) fnd:[MH⁺] 579.2

225. 2,6-Difluoro-N-(4-[6-[3-fluoro-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₀ F₃ N₅ O₅ S (629.66) fnd:[MH⁺] 584.2

226. 6-Fluoro-N-(4-[6-[3-fluoro-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₁ F₂ N₅ O₅ S (611.67) fnd:[MH⁺] 566.2

227. N-(4-[6-[3-Fluoro-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₄ F N₅ O₆ S (623.71) fnd:[MH⁺] 578.1

228. 2,6-Difluoro-N-(4-[6-[3-methyl-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F₂ N₅ O₅ S (625.70) fnd:[MH⁺] 580.2

- 59 -

229. 6-Fluoro-N-(4-{6-[3-methyl-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt
- MS: calc: C₃₁ H₃₄ F N₅ O₅ S (607.71) fnd:[MH⁺] 562.2
230. 2,4-Difluoro-N-(4-{6-[3-methyl-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt
- MS: calc: C₃₁ H₃₃ F₂ N₅ O₅ S (625.70) fnd:[MH⁺] 580.2
231. N-(4-{6-[3-methyl-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt
- MS: calc: C₃₂ H₃₇ N₅ O₆ S (619.75) fnd:[MH⁺] 574.3
232. 3-Fluoro-4-methoxy-N-(4-{6-[3-methyl-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt
- MS: calc: C₃₂ H₃₆ F N₅ O₆ S (637.74) fnd:[MH⁺] 592.2
233. 2,6-Difluoro-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt
- MS: calc: C₃₁ H₃₃ F₂ N₅ O₆ S (641.70) fnd:[MH⁺] 596.1
234. 6-Fluoro-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt
- MS: calc: C₃₁ H₃₄ F N₅ O₆ S (623.71) fnd:[MH⁺] 578.2
235. 2,4-Difluoro-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt
- MS: calc: C₃₁ H₃₃ F₂ N₅ O₆ S (641.70) fnd:[MH⁺] 596.2
236. 4-Methoxy-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

- 60 -

MS: calc: C₃₂ H₃₇ N₅ O₇ S (635.74) fnd:[MH⁺] 590.2

237. 4-Methyl-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₇ N₅ O₆ S (619.75) fnd:[MH⁺] 574.2

238. 2-Fluoro-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₆ F N₅ O₆ S (637.74) fnd:[MH⁺] 592.2

239. 3-Fluoro-4-methoxy-N-(4-{6-[3-methoxy-4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino}-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₆ F N₅ O₇ S (653.74) fnd:[MH⁺] 608.2

240. 2-Fluoro-N-[4-{6-(1H-indol-5-yl)-pyrimidin-4-ylamino}-phenyl]-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₂₈ H₂₂ F N₅ O₄ S (519.56) fnd:[MH⁺] 474.2

241. 2-Fluoro-4-methyl-N-[4-{6-(1-methyl-1H-indol-5-yl)-pyrimidin-4-ylamino}-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₂₇ H₂₄ F N₅ O₄ S (533.59) fnd:[MH⁺] 488.2

242. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-3-fluoro-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₁ F N₆ O₅ S (606.68) fnd:[MH⁺] 561.2

243. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₁ F N₆ O₄ S (590.68) fnd:[MH⁺] 545.2

- 61 -

244. N-(4-[6-[1-(2-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-3-fluoro-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F N₆ O₅ S (620.71) fnd: [MH⁺] 575.2

245. N-(4-[6-[1-(2-Dimethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₁ H₃₃ F N₆ O₄ S (604.71) fnd: [MH⁺] 559.2

246. N-(4-[6-[1-(2-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-3-fluoro-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₇ F N₆ O₅ S (648.76) fnd: [MH⁺] 603.2

247. N-(4-[6-[1-(2-Diethylamino-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₇ F N₆ O₄ S (632.76) fnd: [MH⁺] 587.2

248. 2-Fluoro-4-methyl-N-(4-[6-[1-(2-pyrrolidin-1-yl-ethyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₂ H₃₃ F N₆ O₄ S (616.72) fnd: [MH⁺] 571.2

249. 2-Fluoro-4-methyl-N-(4-[6-[1-(2-pyrrolidin-1-yl-propyl)-1H-indol-5-yl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₅ F N₆ O₄ S (630.75) fnd: [MH⁺] 585.3

250. 2-Fluoro-4-methyl-N-[4-(6-[1-[2-(4-methyl-piperazin-1-yl)-ethyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

MS: calc: C₃₃ H₃₆ F N₇ O₄ S (645.76) fnd: [MH⁺] 600.3

251. 2-Fluoro-4-methyl-N-[4-(6-[1-[2-(4-methyl-piperazin-1-yl)-propyl]-1H-indol-5-yl]-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide formic acid salt

- 62 -

MS: calc: C₃₄ H₃₈ F N₇ O₄ S (659.79) fnd:[MH⁺] 614.3

252. N-(4-{6-[1-(3-Dimethylamino-propyl)-1H-benzoimidazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₂₉ H₃₀ F N₇ O₄ S (591.67) fnd:[MH⁺] 546.3

253. N-(4-{6-[1-(3-Dimethylamino-propyl)-1H-benzoimidazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₂₉ H₂₉ F₂ N₇ O₄ S (609.66) fnd:[MH⁺] 564.3

254. N-(4-{6-[1-(3-Dimethylamino-propyl)-1H-benzoimidazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₂ F N₇ O₄ S (605.70) fnd:[MH⁺] 560.3

255. N-(4-{6-[1-(3-Dimethylamino-propyl)-1H-benzoimidazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₃₀ H₃₃ N₇ O₅ S (603.71) fnd:[MH⁺] 558.3

256. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-benzenesulfonamide formic acid salt

MS: calc: C₂₈ H₂₈ F N₇ O₄ S (577.64) fnd:[MH⁺] 532.3

257. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2,6-difluoro-benzenesulfonamide formic acid salt

MS: calc: C₂₈ H₂₇ F₂ N₇ O₄ S (595.63) fnd:[MH⁺] 550.3

258. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-2-fluoro-4-methyl-benzenesulfonamide formic acid salt

MS: calc: C₂₉ H₃₀ F N₇ O₄ S (591.67) fnd:[MH⁺] 546.3

- 63 -

259. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methyl-benzenesulfonamide formic acid salt


MS: calc: C₂₉ H₃₁ N₇ O₄ S (573.68) fnd:[MH⁺] 528.3

260. N-(4-{6-[1-(2-Dimethylamino-ethyl)-1H-indazol-5-yl]-pyrimidin-4-ylamino}-phenyl)-4-methoxy-benzenesulfonamide formic acid salt

MS: calc: C₂₉ H₃₁ N₇ O₅ S (589.68) fnd:[MH⁺] 544.3


Starting materials and Intermediates:

Example of the synthesis of intermediates according to step 1, step 2, step 3, step 4 and step 7 (compare reaction scheme 1):

A1. 3-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-3-oxo-propionic acid ethylester


A solution of sodium hydride (60%, 12.9 g, 323 mmol) and diethyl carbonate (33 ml, 269 mmol) in absolute toluene (500 ml) is stirred at room temperature for 0.5 h. After stirring under reflux for 10 min a solution of 1-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-ethanone (50 g, 269 mmol) in absolute toluene (250 ml) is slowly added and the reaction mixture is stirred again for 1 h under reflux. The reaction solution is diluted with ice water (500 ml) and neutralized with acetic acid (150 ml). The organic layer is separated, washed with H₂O, dried over MgSO₄, filtered off and concentrated under reduced pressure. Further purification by chromatography [Petrolether/ Ethyl acetate (8:2)] over a silica gel column gives the title compound (72 g) as a colorless solid. TLC, silica gel, glass plates, [Petrolether/ Ethyl acetate (8:2)], R_f = 0.47.

MS: calc.: C₁₄H₁₆O₄ (248.2), fnd: [MH⁺] 249.0

A2. 6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-2-thioxo-2,3-dihydro-1H-pyrimidin-4-one


To a stirred solution of 3-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-3-oxo-propionic acid ethylester (37 g, 150 mmol) and sodium ethylate (20%, 100 ml, 223 mmol) in absolute ethanol (400 ml) thiourea (22.5 g, 298 mmol) is added and the mixture is stirred under reflux for 4 d. After cooling to room temperature the reaction solution is evaporated in vacuo and the resulting residue is dissolved in aqueous 2N HCl (400 ml). The crude product is filtered and washed with H₂O and recrystallized in ethyl acetate and toluene to give the title compound (58 g) as a colorless solid. TLC, silica gel, glass plates, [Toluene/ Acetone (9:1)], R_f = 0.29.

MS: calc.: C₁₃H₁₂N₂O₂S (260.2), fnd: [MH⁺] 261.0

A3. 6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-3H-pyrimidin-4-one


6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-2-thioxo-2,3-dihydro-1H-pyrimidin-4-one (30 g, 115 mmol) is dissolved in H₂O (300 ml). After addition of an aqueous solution of ammonia (25%, 130 ml) and Raney-Nickel WII (30 g) in H₂O (20 ml) the reaction mixture is stirred under reflux for 16 h. The reaction mixture is filtered from the solid material, which is washed again with a mixture of ammonia and H₂O (1:2, 300 ml). The combined aqueous layers are evaporated in the vacuo and coevaporated with toluene (3 x) to give the title compound (31 g) as a colorless solid. TLC, silica gel, glass plates, [CH₂Cl₂/ MeOH (95:5)], R_f = 0.36.

MS: calc: C₁₃H₁₂N₂O₂ (228.2), fnd: [MH⁺] 229.0

A4. 4-Chloro-6-(2-methyl-2,3-dihydro-benzofuran-5-yl)-pyrimidine

6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-3H-pyrimidin-4-one (31 g, 136 mmol) is dissolved in POCl₃ (120 ml) and stirred under reflux for 2 h. After cooling to room temperature the reaction solution is poured on ice water (1.5 l) and stirred for 1 h. The solution is neutralized by the addition of K₂CO₃ powder and extracted with ethyl acetate (3 x 500 ml). The organic phase is dried over MgSO₄, filtered off and concentrated under reduced pressure. Further purification by chromatography [Toluene/ Acetone (95:5)] over a silica gel column gives the title compound (24 g) as a beige solid. TLC, silica gel, glass plates [Toluene/ Acetone (9:1)], R_f = 0.43.

MS: calc: C₁₃H₁₁ClN₂O (246.7), found: [MH⁺] 247.2, 249.2


A5. {4-[6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-pyrimidin-4-yl-amino]-phenyl}-carbamic acid tert-butyl ester

- 66 -

4-Chloro-6-(2-methyl-2,3-dihydro-benzofuran-5-yl)-pyrimidin (10 g, 41 mmol) is dissolved in a solution of Et_3N (9 ml) and absolute DMF (50 ml), (4-Amino-phenyl)-carbamic acid tert-butylester (9.3 g, 45 mmol) is added and the reaction mixture is stirred for 2 h at 140°C . The reaction solution is diluted with CH_2Cl_2 (80 ml) and extracted with a semisaturated aqueous NaCl solution (80 ml). The organic phase is dried over MgSO_4 , filtered off and concentrated under reduced pressure. Further purification by chromatography [$\text{CH}_2\text{Cl}_2/\text{MeOH}$ (95:5)] over a silica gel column gives the title compound (8.2 g) as a colorless solid. TLC, silica gel, glass plates, [$\text{CH}_2\text{Cl}_2/\text{MeOH}$ (95:5)], $R_f = 0.35$.

MS: calc: $\text{C}_{24}\text{H}_{28}\text{N}_4\text{O}_3$ (318.5), fnd: $[\text{MH}^+ - \text{Boc}]$ 319.1; $[\text{MH}^+ - 56]$ 363.2; $[\text{MH}^+]$ 419.1; $[2\text{MH}^+]$ 836.3

A6. N-[6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-pyrimidin-4-yl]-benzene-1,4-diamine dihydrochloride




A suspension of {4-[6-(2-Methyl-2,3-dihydro-benzofuran-5-yl)-pyrimidin-4-yl-amino]-phenyl}-carbamic acid *tert*-butyl ester (19.6 g; 47 mmol) in dioxane (90 ml) is admixed with a saturated solution of HCl in dioxane (60 ml) and stirred at room temperature for 2 h. The reaction mixture is diluted with diethyl ether and the resulting precipitate is filtered off under an N_2 atmosphere and washed with diethyl ether (3 x 50 ml). Drying under reduced pressure gives the title compound (16.5 g) as a colorless solid.

MS: calc: $\text{C}_{19}\text{H}_{18}\text{N}_4\text{O}$ (318.4), fnd: $[\text{MH}^+]$ 319.3

Example of the synthesis of intermediates according to step 5 and step 6 (compare reaction scheme 1):

A7. [4-(6-Chloro-pyrimidin-4-ylamino)-phenyl]-carbamic acid tert-butyl ester




A solution of 4,6-dichloropyrimidine (53.6 g), (4-amino-phenyl)-carbamic acid tert-butyl ester (50.0 g), DABCO (2.7 g) and diisopropyl ethyl amine (46.6 g) in dimethylformamide is stirred at 120°C under an atmosphere of nitrogen for 2 h. The solvent is removed in vacuo. The residue is dissolved in ethyl acetate and washed with 1M citric acid and saturated aqueous NaCl solution. The organic layer is dried over MgSO_4 and filtered through a plug of neutral alumina. The filtrate is concentrated in vacuo. 48.2 g of pure product is obtained after crystallization from *tert*-BuOMe and hexane as off-white solid.

- 67 -

$^1\text{H-NMR}$ (DMSO- d_6)**8** (ppm): 9.71 (s, 1H, -NH); 9.29 (s, 1H, -NH); 8.41 (d, $J = 0.6$ Hz, 1H); 7.44 (s, 4H); 6.69 (d, $J = 0.6$ Hz, 1H); 1.47 (s, 9H).

MS (ESI): $m/e = 321.0$ (MH^+ 100%); 265.2 ($\text{MH}^+ - 56$, 62%); 221.3 ($\text{MH}^+ - 100$, 12%).

A8. {4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl}-carbamic acid tert -butyl ester




[4-(6-Chloro-pyrimidin-4-ylamino)-phenyl]-carbamic acid tert-butyl ester (7.0 g) and 4-fluorophenyl-boronic acid (4.6 g) is dissolved in degassed dioxane. After addition of 2 M aqueous Na_2CO_3 solution (33 ml) and trans-dichlorobis(tricyclohexylphosphine)palladium (0.8 g) the reaction mixture is stirred at 90°C under an atmosphere of nitrogen for 3 h. Saturated aqueous NaCl solution is added, the organic layer is separated, dried over MgSO_4 and filtered through a plug of neutral alumina. The filtrate is concentrated in vacuo. 5.2 g of pure product is obtained after crystallization from ethyl acetate as off-white solid.

$^1\text{H-NMR}$ (DMSO- d_6)**8** (ppm): 9.52 (s, 1H, -NH); 9.26 (s, 1H, -NH); 8.65 (d, $J = 0.7$ Hz, 1H); 8.07 (m, 2H); 7.55 (d, $J = 9.0$ Hz, 2H); 7.43 (d, $J = 9.0$ Hz, 2H); 7.35 (dxd, $J_1 = J_2 = 8.9$ Hz, 2H); 7.13 (d, $J = 0.7$ Hz, 1H); 1.48 (s, 9H).

MS(ESI): $m/e = 381.1$ (MH^+ , 100%); 325.2 ($\text{MH}^+ - 56$, 59%), 281.3 ($\text{MH}^+ - 100$, 18%).

Example of the synthesis according to step 7 and step 8 (compare reaction scheme 1):

A9. N-[6-(4-Fluoro-phenyl)-pyrimidin-4-yl]-benzene-1,4-diamine



{4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl}-carbamic acid tert -butyl ester (5.2 g) is dissolved in dioxane. 4N HCl in dioxane is added and the well stirred reaction mixture is heated to 50°C for several hours. The HCl salt of the product is precipitated by addition of Et_2O . The precipitate is filtered and dis-


- 68 -

tributed between aqueous Na_2CO_3 and ethyl acetate. The organic layer is separated and dried over MgSO_4 . The solvent is removed in vacuo. 4.9 g of pure product is obtained after crystallization from ethyl acetate and hexane as off-white solid.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm): 9.17 (s, 1H, -NH); 8.55 (s, 1H); 8.02 (m, 2H); 7.32 (dxd, $J_1 = J_2 = 8.8$ Hz, 2H); 7.21 (d, 2H); 6.98 (s, 1H); 6.60 (d, $J = 8.8$ Hz, 2H); 4.95 (s, 2H, $-\text{NH}_2$).

MS(ESI). $m/e = 281.3$ (MH^+ , 100%).


1. **2,6-Difluoro-N-[4-[6-(4-fluoro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide**



Synthesis details are described in the section Final Products.

Example of the synthesis according to step 9, step 10 and step 11:

A10. **N-(6-Chloro-pyrimidin-4-yl)-benzene-1,4-diamine**




[4-(6-Chloro-pyrimidin-4-ylamino)-phenyl]-carbamic acid tert-butyl ester (160.4 g) is dissolved in dioxane and treated with 4N HCl in dioxane at 50°C over night under an atmosphere of nitrogen. The hydrochloride of the product is precipitated with Et_2O and isolated by filtration. Precipitation from the aqueous solution of the collected hydrochloride with 2M aqueous Na_2CO_3 solution yields 110.0 g of the free base as off-white solid.

$^1\text{H-NMR}$ (DMSO-d_6) δ (ppm): 9.40 (s, 1H, -NH); 8.31 (s, 1H); 7.11 (d, $J = 8.3$ Hz, 2H); 6.58 (s, 1H), 6.54 (d, $J = 8.3$ Hz, 2H); 5.01 (s, 2H, $-\text{NH}_2$).

MS (ESI): $m/e = 221.2$ (MH^+ , 100%).

A11. N-[4-(6-chloro-pyrimidin-4-ylamino)-phenyl]-2,4-difluoro-benzenesulfonamide




N-(6-Chloro-pyrimidin-4-yl)-benzene-1,4-diamine (11.1 g) and 2,6-lutidine (6.5 g) is dissolved in dimethoxyethane. After slow addition of 2,4-difluoro-benzenesulfonyl chloride (11.7 g) the reaction mixture is stirred under an atmosphere of nitrogen over night at ambient temperature. After addition of 1N HCl and saturated aqueous NaCl solution the organic layer is separated and concentrated in vacuo. 17.7 g of pure product is obtained after crystallization from methanol / water as off-white solid.

¹H-NMR (DMSO-d₆) δ (ppm): 10.50 (s, 1H, -NH); 9.78 (s, 1H, -NH); 8.42 (s, 1H); 7.86 (dxdxd, J₁ = J₂ = 8.6 Hz, J₃ = 6.4 Hz); 7.52 (d, J = 8.8 Hz, 2H, and m, 1H); 7.24 (dxdxd, J₁ = J₂ = 8.6 Hz, J₃ = 1.7 Hz); 7.09 (d, J = 8.8 Hz, 2H); 6.73 (s, 1H).

MS (ESI): m/e = 397.0 (MH⁺, 100%).


2. 2,4-Difluoro-N-(4-[6-[4-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-pyrimidin-4-ylamino]-phenyl)-benzenesulfonamide




Synthesis details are described in the section Final Products.

- 70 -

Example for the synthesis of a boronic acid ester of formula 9b - Reaction scheme 2:



A12. Methanesulfonic acid 1-formyl-piperidin-4-ylmethyl ester




To a well stirred solution of piperidin-4-yl-methanol (69.0 g) in dichloromethane methyl formate (46.8 g) is added with care. The reaction is stirred at ambient temperature for 6 h. The solvent is completely removed in vacuo. The residual pale yellow oil is redissolved in dichloromethane and triethylamine (106.5 g) is added to the well stirred solution. A solution of methanesulfonyl chloride (82.5 g) is added dropwise at 0°C and after complete addition the reaction mixture is stirred at ambient temperature for additional 4 h. The organic layer is extracted with 0.5 N aqueous HCl and dried over MgSO₄. The solvent is removed in vacuo. Pure product 69.9 g is obtained as off-white solid after crystallization from cyclohexane.

MS: calc: C₈H₁₅NO₄S (221.28)

find: [MH⁺]: 222.1

- 71 -


A13. 4-[4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenoxy-methyl]-piperidine-1-carbaldehyde



Sodium hydride (60% dispersion in oil, 1.1 g) is washed with hexane and suspended in dry dimethylformamide. 4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenol (5.2 g) is added in small portions to the well stirred suspension under cooling to 0°C and stirring is continued under an atmosphere of nitrogen for 1 h. Methanesulfonic acid 1-formyl-piperidin-4-ylmethyl ester (6.1 g) is added and the reaction mixture is heated to 100°C for 1 h. After cooling to ambient temperature the reaction is quenched with water and the product is extracted into ethyl acetate. After drying over MgSO₄ the solvent is removed in vacuo. The residue is crystallized from acetonitrile to yield 4.4 g of pure product as an off-white solid.

¹H-NMR(CDCl₃) δ(ppm): 8.0 (s, 1H); 7.7 (d, 2H); 6.8 (d, 2H); 4.5 (m, 1H); 3.8 (t, 2H); 3.6 (m, 1H); 2.0 (m, 2H + 1H); 1.3 (s, 12H); 1.2 (m, 2H).

A14. 1-Methyl-4-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenoxy-methyl]-piperidine




Borane (1M solution in tetrahydrofuran, 25.0 ml) is dropwise added to a solution of 4-[4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenoxy-methyl]-piperidine-1-carbaldehyde (4.3 g) in tetrahydrofuran, then the stirred mixture is heated to gentle reflux for 1.5 h under an atmosphere of nitrogen. After addition of further borane solution (2.5 ml) the mixture is refluxed for another 5 h before addition of pinacol (5.9 g) and again refluxing for 2 h. The volatile materials are removed in vacuo. The residue is purified by Kugelrohr distillation to deliver 1.5 g of the pure product (bp. 160°C, 0.1 mbar) after a forerun of pinacol as colourless oil, which solidified on standing.


¹H-NMR(CDCl₃) δ(ppm): 7.7 (d, 2H); 6.8 (d, 2H); 3.7 (t, 2H); 2.8 (m, 2H); 2.2 (s, 3H); 1.7 (m, 2H + 2H + 1H); 1.4 (m, 2H); 1.3 (s, 12H).

- 72 -

Additional example for the synthesis of a boronic acid ester of formula 9b— reaction scheme 3:



A15. [2-(5-Iodo-indol-1-yl)-ethyl]-dimethyl-amine hydrochloride




Oil free sodium hydride (prepared from 4.0g 60% dispersion in oil by washing with hexane) is suspended in DME, DMSO (9 : 1). To the well-stirred suspension 5-Iodo-1H-indole (9.72 g) is added in portions. After stirring for 30 min (2-chloro-ethyl)-dimethyl-amine hydrochloride (6.91g) is added and the reaction mixture is stirred at 70°C for 16 h under an atmosphere of nitrogen. The reaction mixture is quenched by slow addition of ice-cold water. After saturation of the aqueous layer with solid NaCl the organic layer is separated and concentrated in vacuo. The aqueous layer is extracted with AcOEt. All aqueous layers are combined, washed with brine and dried over MgSO₄. After evaporation the crude product is chromatographed on neutral alumina (act. 2-3) using AcOEt as eluent. The title compound (12.54g) is obtained as a pale yellow oil.

MS: calc: C₁₂H₁₅IN₂ (314.17) fnd:[MH⁺]: 315.1

The hydrochloride is obtained as an off-white solid by dissolving the free base in Et₂O and slow addition of a small excess of 4N HCl in dioxane.

A16. Dimethyl-[2-[5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-indol-1-yl]-ethyl]-amine



[2-(5-Iodo-indol-1-yl)-ethyl]-dimethyl-amine hydrochloride (10.0 g) is suspended in dry dioxane. After addition of dry Et₃N (14.42 g), PdCl₂(dppf)·CH₂Cl₂ (0.35g) and 4,4,5,5-Tetramethyl-1,3,2-dioxaborolane (5.47

- 73 -

g) the reaction mixture is stirred at 80 °C for about 1 h under an atmosphere of nitrogen. After addition of ice cold water the reaction mixture is concentrated in vacuo. The residue is extracted with several portions AcOEt. The combined organic layers are dried over MgSO₄. After filtration the solvent is removed in vacuo. The crude product is filtered through a plug of neutral alumina (act. 2-3) using AcOEt as eluent and further purified by bulb to bulb distillation (200 °C – 220 °C; 1.8×10^{-2} mbar). The title compound (8.90g) is obtained as a pale yellow oil.

MS: calc: C₁₈H₂₇BN₂O₂ (314.24) fnd:[MH⁺]: 315.2

¹H-NMR (DMSO-d₆) δ (ppm): 7.94 (s, 1H); 7.45 (s, 2H); 7.37 (d, J = 3.2 Hz, 1H); 6.46 (d, J = 3.2 Hz, 1H); 4.25 (t, J = 6.6 Hz, 2H); 2.60 (t, J = 6.6 Hz, 2H); 2.18 (s, 6H); 1.31 (s, 12H).

Biological assays for the investigation of T cell associated kinase inhibitors

The compounds of formula 1 in free form or in pharmaceutically acceptable salt form exhibit valuable pharmacological properties, e.g. inhibition of protein kinases like p90 ribosomal S6 kinase (Rsk) family, Src family kinases, e.g. Lck or Protein Kinase C (PKC), e.g. PKC isoforms like α or θ activity, inhibition of T lymphocyte activation and proliferation, e.g. by inhibiting production of cytokines by T lymphocytes, e.g. IL-2, by inhibiting the proliferative response of T lymphocytes to cytokines, e.g. IL-2, as indicated in in vitro tests and are therefore indicated for therapy.

A. Biochemical Tests**1. Protein Kinase C assay**

The compounds of formula 1 are tested for their activity on different protein kinases and PKC isoforms using a scintillation proximity assay (SPA, Amersham International plc). ^{33}P -labeled peptides are captured onto streptavidin coated yttrium silicate SPA beads. β -particles, emitted from the captured ^{33}P -labeled substrate in close proximity to the bead are able to excite the scintillant, resulting in the generation of quantifiable light. The assay is performed in a 96-well polystyrene microtiterplate (1450-514, Isoplates, Wallac, Turku, Finland). The reaction mixture (50 μl) contains 10 μl of the test compound together with 10 μl of the relevant kinase, diluted in the relevant dilution buffer, 5 μl of 10 μM phorbol myristate acetate (PMA) in H_2O , 5 μl of 1,6 mM phosphatidylserine (Sigma-Aldrich Chemie GmbH, Steinheim, Germany) in 20 mM Tris/HCL buffer pH 7.4, 5 μl of 0,3 % BSA in H_2O , 5 μl of 30 μM relevant substrate and 10 μl of 5 μM ATP and 0,1 μCi of ^{33}P -ATP (Amersham, Freiburg, Germany) in 200 mM Tris/HCL pH 7.5 and 200 mM MgCl_2 . Incubation is performed for 40 min at room temperature (RT) without shaking. Reaction is stopped by adding 150 μl of cold stop solution containing 10 mM ATP, 5 mM EGTA pH 7.5, 0,1 % Triton X-100 and 0,2 mg streptavidine coated yttrium silicate SPA beads (Amersham, RPNQ 0012). The sealed plate is incubated for 60 min at RT. Thereafter the MTP is counted in a Microbeta Jet (Wallac). IC_{50} measurement is performed on a routine basis by incubation a serial dilution of inhibitor at concentrations ranging between 0.01 and 100 μM according to the method described above. Background values are the signals of the reaction mixture without addition of the relevant kinase and are subtracted from all values. 100 % values are the signals of the reaction mixture without addition of inhibitors.

Dependent on the efficacy of the inhibitors in the various test systems, corresponding IC_{50} values are calculated from concentration-inhibition curves by nonlinear regression analysis using the program GraphPad Prism (GraphPad Software Inc., San Diego, CA).

2. Protein Kinase C α Assay

- 75 -

Human recombinant PKC α was obtained from Panvera (Invitrogen GmbH, Karlsruhe, Germany) and is used under the assay conditions as described above (Section A.1). The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The assay additionally contains 0.5 mM CaCl_2 . The kinase reaction is performed with the biotinylated PKC α pseudosubstrate solved in H_2O . The examples 16, 37, 119, 120 and 150 inhibit PKC α in this assay with an IC_{50} between 1 and 18 μM .

3. Protein Kinase C β 1 Assay

Human recombinant PKC β 1 was obtained from Panvera and is used under the assay conditions as described above (Section A.1). The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The assay additionally contains 0.5 mM CaCl_2 . The kinase reaction is performed with the biotinylated PKC α pseudosubstrate solved in H_2O . The examples 29, 45, 52, 128 and 139 inhibit PKC β 1 in this assay with an IC_{50} between 2 and 40 μM .

4. Protein Kinase C δ Assay

Human recombinant PKC δ was obtained from Panvera and is used under the assay conditions as described above (Section A.1). The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The kinase reaction is performed with the biotinylated PKC η pseudosubstrate solved in H_2O . The examples 122, 132, 138 and 168 inhibit PKC δ in this assay with an IC_{50} between 2 and 18 μM .

5. Protein Kinase C ϵ Assay

Human recombinant PKC ϵ was obtained from Panvera and is used under the assay conditions as described above (Section A.1). The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The kinase reaction is performed with the biotinylated PKC η pseudosubstrate solved in H_2O . The examples 27, 38, 47 and 117 inhibit PKC ϵ in this assay with an IC_{50} between 4 and 21 μM .

6. Protein Kinase C η Assay

Human recombinant nickel bead purified PKC η from Sf21 insect cells is used under the assay conditions as described above (Section A.1). The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The kinase reaction is performed with the biotinylated PKC η pseudosubstrate solved in H_2O . The examples 52, 55, 118 and 135 inhibit PKC η in this assay with an IC_{50} between 6 and 50 μM .

7. Protein Kinase C θ Assay

Human recombinant nickel bead purified PKC θ from Sf21 insect cells is used under the assay conditions as described above (Section A.1). The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes

- 76 -

pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The kinase reaction is performed with the biotinylated PKC θ pseudosubstrate solved in H₂O. The examples 8, 22, 46, 115, 118 and 161 inhibit PKC θ in this assay with an IC₅₀ < 1 μ M.

8. Protein Kinase C ι Assay

Human recombinant nickel bead purified PKC ι from Sf21 insect cells is used under the assay conditions as described above (Section A.1) but in the absence of PMA and phosphatidylserine. The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The kinase reaction is performed with the biotinylated PKC ι / ζ pseudosubstrate solved in H₂O. The examples 145, 162 and 173 inhibit PKC ι in this assay with an IC₅₀ between 7 and 34 μ M.

9. Protein Kinase C ζ Assay

Human recombinant PKC ζ was obtained from Panvera and is used under the assay conditions as described above (Section A.1) but in the absence of PMA and phosphatidylserine. The enzyme is diluted in PKC dilution buffer containing 1 mM Hepes pH 7.4, 0.5 mM DTT and 0.001 % Triton X-100. The kinase reaction is performed with the biotinylated PKC ι / ζ pseudosubstrate solved in H₂O. The examples 143 and 175 inhibit PKC ζ in this assay with an IC₅₀ between 15 and 25 μ M.

10. Protein Kinase A Assay

Human recombinant PKA was obtained from Panvera and is used in a scintillation proximity assay described in Section A.1. The reaction mixture (50 μ l) contains 15 μ l of the test compound together with 10 μ l of PKA in 1 mM Hepes pH 7.4, 0.001 % Triton X-100 and 1 mM DTT, 5 μ l of 0.3 % BSA in H₂O, 10 μ l of 15 μ M biotinylated PKA substrate peptide (Upstate) and 10 μ l of 5 μ M ATP and 0.1 μ Ci of ³³P-ATP in 200 mM Tris/HCL pH 7.5 and 100 mM MgCl₂. Incubation is performed for 40 min at room temperature (RT) without shaking. Reaction is stopped by adding 150 μ l of cold stop solution containing 10 mM ATP, 5 mM EGTA pH 7.5, 0.1 % Triton X-100 and 0.2 mg streptavidine coated yttrium silicate SPA beads (Amersham, RPNQ 0012). The sealed plate is incubated for 60 min at RT. Thereafter the MTP is counted in a Microbeta Jet (Wallac). IC₅₀ measurement is performed on a routine basis by incubation a serial dilution of inhibitor at concentrations ranging between 0.01 and 100 μ M according to the method described above. Background values are the signals of the reaction mixture without addition of the relevant kinase and are subtracted from all values. 100 % values are the signals of the reaction mixture without addition of inhibitors. IC₅₀ values are calculated from the graph by sigmoidal curve fitting. The examples 10, 17, 117 and 121 inhibit PKA in this assay with an IC₅₀ between 3 and 16 μ M.

11. Lck (p56^{Lck}) Assay

Human recombinant Lck was obtained from Upstate (Dundee, UK) and is used in a scintillation proximity assay described in Section A.1. The reaction mixture (50 μ l) contains 15 μ l of the test compound together with 10 μ l of Lck in 1 mM Hepes pH 7.4, 0.001 % Triton X-100 and 1 mM DTT, 5 μ l of 0.3 % BSA in H₂O,

- 77 -

10 μ l of 150 μ M biotinylated synthetic peptide obtained from Biotrend (Köln, Germany) and 10 μ l of 5 μ M ATP and 0,1 μ Ci of 33 P-ATP in 200 mM Tris/HCL pH 7.5 and 100 mM MgCl₂. Incubation is performed for 40 min at room temperature (RT) without shaking. Reaction is stopped by adding 150 μ l of cold stop solution containing 10 mM ATP, 5 mM EGTA pH 7.5, 0,1 % Triton X-100 and 0,2 mg streptavidine coated yttrium silicate SPA beads (Amersham, RPNQ 0012). The sealed plate is incubated for 60 min at RT. Thereafter the MTP is counted in a Microbeta Jet (Wallac). IC₅₀ measurement is performed on a routine basis by incubation a serial dilution of inhibitor at concentrations ranging between 0,01 and 100 μ M according to the method described above. Background values are the signals of the reaction mixture without addition of the relevant kinase and are subtracted from all values. 100 % values are the signals of the reaction mixture without addition of inhibitors. IC₅₀ values are calculated from the graph by sigmoidal curve fitting. The examples 128 and 131 inhibit Lck in this assay with an IC₅₀ < 3 μ M.

12. Rsk1, Rsk2, Rsk3 Assay

To investigate the effect of a test compound on p90 ribosomal S6 kinase (Rsk) family, the compounds are externally tested in KinaseProfiler assay (Upstate Ltd, Dundee, UK) described in product guide brochure. In brief, in a final reaction volume of 25 μ l, 5 – 10 mU of human recombinant Rsk are incubated with 8 mM MOPS pH 7.0, 0.2 mM EDTA, 30 μ M KKKNRTLSVA, 10 mM MgAcetate and 10 μ M [γ - 33 P-ATP]. The reaction is initiated by addition of the MgATP mix. After incubation for 40 minutes at RT, the reaction is stopped by the addition of 5 μ l of a 3 % phosphoric acid solution. 10 μ l of the reaction is then spotted onto P30 filtermat and washed three times for 5 minutes in 75 mM phosphoric acid and once in methanol prior to drying and scintillation counting. The IC₅₀ value is estimated by preparing a 10 point curve using a $\frac{1}{2}$ log dilution series (IC₅₀Profiler, Upstate). The examples 129, 131 and 173 inhibit Rsk1 in this assay with an IC₅₀ < 1 μ M.

B Cellular Tests

1. CD3/CD28 costimulation assay

The assay is performed with freshly isolated primary human CD4⁺ T lymphocytes. CD4⁺ T lymphocytes from whole blood are prepared using negative selection as previously described (Hatzelmann and Schudt, J. Pharmacol. Exp. Ther. 2001; 297: 267-279). In brief, peripheral blood mononuclear cells (PBMC) are isolated by density gradient centrifugation using a Percoll gradient (ρ = 1.077 g/ml). 1×10^7 PBMC were then resuspended in 30 μ l of PBS containing 0.5 % FCS and human CD4⁺ T lymphocytes are isolated by depletion of non CD4⁺ T cells. For this, non-CD4⁺ T cells are indirectly magnetically labeled with 10 μ l of a cocktail of biotin-conjugated monoclonal antibodies (against CD8, CD14, CD16, CD19, CD36, CD56, CD123, TCR $\gamma\delta$; MACS CD4⁺ T cell isolation kit II, Miltenyi Biotec), as primary labeling reagent, and anti-biotin monoclonal antibodies conjugated to MicroBeads, as secondary labeling reagent. The magnetically labeled non-CD4⁺ T cells are depleted by retaining them on a MACS column in the magnetic field of a MACS separator (Miltenyi Biotec, Bergisch Gladbach, Germany) according to the manufacturer's instruc-

- 78 -

tion. CD4+ T cells are resuspended in RPMI 1640 containing 10 % heat-inactivated FCS, 2 mM L-glutamine, 100 U/ml penicillin and 100 µg/ml streptomycin (Gibco Life Technologies).

2×10^5 CD4+ T cells in a total assay volume of 200 µl are stimulated via the T-cell receptor and the costimulatory molecule CD28 by using corresponding selective mAbs as previously described (Hatzelmann and Schudt, J. Pharmacol. Exp. Ther. 2001; 297: 267-279). The assay is performed in 96 well tissue culture plates (655180, flat bottom, Greiner, Frickenhausen, Germany). Cells were maintained at 37°C in a humidified atmosphere of 5 % CO₂ in an incubator (type BB6220 CU, Heraeus Instruments, Hanau, Germany). For determination of IL-2 level, all assays are performed in duplicate, and after 48 h of growth, supernatants are removed, pooled in 96 well plates (650101, U-shape, Greiner) and stored at -20°C before measurement of IL-2 with a commercially available enzymimmunoassay kit from Coulter-Immunotech Diagnostics (Marseille, France) according to the manufacturer's instruction. For each experiment the appropriate dilution factor for IL-2 is determined. Dilutions are performed in diluent D (Coulter-Immunotech Diagnostics) and IL-2 for one condition is determined from the pool fraction in duplicate in a ELISA-reader (Rainbow, Tecan, Crailsheim, Germany) at 450 nm.

To investigate the effect of a test compound on the IL-2 release of stimulated CD4+ T cells, six three-fold dilution steps in duplicates per test compound are performed (final DMSO concentration 0,1 %). Low control values are the signals from non stimulated CD4+ T cells; high controls are the signals from stimulated CD4+ T cells without any test samples. Low controls are subtracted from all values. The inhibition obtained in the presence of a test compound is calculated as percent inhibition of the high control. The concentration of test compounds resulting in 50 % inhibition (IC₅₀) is determined from the dose-response curves. The examples 13, 19, 20 and 136 inhibit IL-2 release from stimulated CD4+ T lymphocytes in this assay with an IC₅₀ between 1 and 9 µM.

2. Mixed Lymphocyte Reaction (MLR)

For MLR cultures 4×10^5 responder T cells are incubated in duplicates with 2×10^5 mitomycin C-treated allogeneic stimulator T cells in a total volume of 200 µl RPMI 1640 medium supplemented with 10 % FCS, 2 mM L-glutamine, 100 U/ml penicillin and 100 µg/ml streptomycin (Gibco Life Technologies) in 96 well tissue culture plates (655180, flat bottom, Greiner). Cells are maintained at 37°C in a humidified atmosphere of 5 % CO₂. After 48 h of growth, cells are pulsed with 1 µCi [³H]thymidine and incorporation of [³H]thymidine is measured with a Topcount radioactive counter (Packard). To investigate the effect of a test compound on MLR-mediated T cell proliferation six three-fold dilution steps in duplicates per test compound are performed (final DMSO concentration 0,1 %). Low control values are the proliferation of responder cells alone; high controls are from mixed lymphocyte cells without any test samples. Low controls are subtracted from all values. The inhibition obtained in the presence of a test compound is calculated as percent inhibition of the high control. The concentration of test compounds resulting in 50 %

- 79 -

inhibition (IC_{50}) is determined from the dose-response curves. The examples 91, 128, 133 and 137 inhibit MLR-mediated T-cell proliferation in this assay with an $IC_{50} < 1 \mu M$.

The compounds according to the invention are, therefore, useful in the treatment and/or prevention of diseases or disorders mediated by T lymphocytes and/or PKC, e.g. acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock. The compounds according to the invention are also useful in the treatment and/or prevention of T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitis, seborrheic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

The invention further relates to a method for the treatment of mammals, including humans, which are suffering from one of the abovementioned illnesses. The method is characterized in that a therapeutically active and pharmacologically effective and tolerable amount of one or more of the compounds according to the invention is administered to the ill mammal.

The invention further relates to the compounds according to the invention for use in the treatment and/or prophylaxis of illnesses, especially the illnesses mentioned.

The invention also relates to the use of the compounds according to the invention for the production of pharmaceutical compositions which are employed for the treatment and/or prophylaxis of the illnesses mentioned.

The invention furthermore relates to pharmaceutical compositions for the treatment and/or prophylaxis of the illnesses mentioned, which contain one or more of the compounds according to the invention.

The pharmaceutical compositions are prepared by processes, which are known per se and familiar to the person skilled in the art. As pharmaceutical compositions, the compounds according to the invention (= active compounds) are either employed as such, or preferably in combination with suitable pharma-

- 80 -

ceutical auxiliaries and/or excipients, e.g. in the form of tablets, coated tablets, capsules, caplets, suppositories, emulsions, suspensions, gels or solutions, the active compound content advantageously being between 0.1 and 95% and where, by the appropriate choice of the auxiliaries and/or excipients, a pharmaceutical administration form (e.g. a delayed release form or an enteric form) exactly suited to the active compound and/or to the desired onset of action can be achieved.

The person skilled in the art is familiar with auxiliaries or excipients which are suitable for the desired pharmaceutical formulations on account of his/her expert knowledge. In addition to solvents, gel formers, ointment bases and other active compound excipients, for example antioxidants, dispersants, emulsifiers, preservatives, solubilizers, colorants, complexing agents or permeation promoters, can be used.

The administration of the pharmaceutical compositions according to the invention may be performed in any of the generally accepted modes of administration available in the art. Illustrative examples of suitable modes of administration include intravenous, oral, nasal, parenteral, topical, transdermal and rectal delivery. Oral and intravenous delivery is preferred.

For the treatment of disorders of the respiratory tract, the compounds according to the invention are preferably also administered by inhalation in the form of an aerosol; the aerosol particles of solid, liquid or mixed composition preferably having a diameter of 0.5 to 10 μm , advantageously of 2 to 6 μm .

Aerosol generation can be carried out, for example, by pressure-driven jet atomizers or ultrasonic atomizers, but advantageously by propellant-driven metered aerosols or propellant-free administration of micronized active compounds from inhalation capsules.

Depending on the inhaler system used, in addition to the active compounds the administration forms additionally contain the required excipients, such as, for example, propellants (e.g. Frigen in the case of metered aerosols), surface-active substances, emulsifiers, stabilizers, preservatives, flavorings, fillers (e.g. lactose in the case of powder inhalers) or, if appropriate, further active compounds.


For the purposes of inhalation, a large number of apparatuses are available with which aerosols of optimum particle size can be generated and administered, using an inhalation technique which is as right as possible for the patient. In addition to the use of adaptors (spacers, expanders) and pear-shaped containers (e.g. Nebulator®, Volumatic®), and automatic devices emitting a puffer spray (Autohaler®), for metered aerosols, in particular in the case of powder inhalers, a number of technical solutions are available (e.g. Diskhaler®, Rotadisk®, Turbohaler® or the inhaler described in European Patent Application EP 0 505 321), using which an optimal administration of active compound can be achieved.

- 81 -

For the treatment of dermatoses, the compounds according to the invention are in particular administered in the form of those pharmaceutical compositions which are suitable for topical application. For the production of the pharmaceutical compositions, the compounds according to the invention (= active compounds) are preferably mixed with suitable pharmaceutical auxiliaries and further processed to give suitable pharmaceutical formulations. Suitable pharmaceutical formulations are, for example, powders, emulsions, suspensions, sprays, oils, ointments, fatty ointments, creams, pastes, gels or solutions.


The pharmaceutical compositions according to the invention are prepared by processes known per se. The dosage of the active compounds is carried out in the order of magnitude customary for kinase inhibitors. Topical application forms (such as ointments) for the treatment of dermatoses thus contain the active compounds in a concentration of, for example, 0.1-99%. The dose for administration by inhalation is customarily between 0.1 and 10 mg per day. The customary dose in the case of systemic therapy (p.o.) is between 0.3 and 30 mg/kg per day, (i. v.) is between 0.3 and 30 mg/kg/h.

- 82 -

Patent claims**1. Compounds of formula 1**

in which

- R1 is phenyl,
 phenyl substituted by R3 and/or R4,
 naphthalenyl,
 naphthalenyl substituted by R5 and/or R6,
 aryl1,
 aryl1 substituted by R7 and/or R8,
 R9,
 R10 or
 R11,
 R2 is phenyl,
 phenyl substituted by R12 and/or R13,
 naphthalenyl,
 naphthalenyl substituted by R14 and/or R15,
 aryl2,
 aryl2 substituted by R16 and/or R17
 or a radical selected from



- R3 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl,

- 83 -

aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$ wherein

R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein

R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

n is an integer from 0 to 4,

m is an integer from 2 to 4,

R4 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R5 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$,

R6 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,


Aryl1 furanyl, thiophenyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl or dibenzofuranyl,

R7 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzy-

- 84 -

loxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R_{31})R_{32}$, $-CH_2CH(OH)CH_2N(R_{31})R_{32}$ or $-O-(CH_2)_m-N(R_{31})R_{32}$,


- R8 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R9 is unsubstituted pyrrolyl, pyrazolyl, imidazolyl, indolyl, indazolyl, benzimidazolyl or benztriazolyl, or a radical selected from



wherein

- R91 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R_{31})R_{32}$, $-CH_2CH(OH)CH_2N(R_{31})R_{32}$ or $-O-(CH_2)_m-N(R_{31})R_{32}$,
- R92 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R10 is a radical selected from

- 85 -




wherein

R101 is 1-4C-alkyl, 2,2,2-trifluoroethyl or 3,3,3-trifluoropropyl,

R102 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R11 is a radical selected from



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R112)R113$ or $-CH_2CH(OH)CH_2N(R112)R113$, wherein

R112 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R113 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R112 and R113 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

- 86 -

- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzo-furanyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,
- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds with the proviso that the following compounds are excluded

4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

N-[4-[6-(2-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

N-[4-[6-(4-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

N-(3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl)-acetamide,

N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,

4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,

N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,

- 87 -

N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide, and
 4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

2. Compounds of formula 1 according to claim 1 in which

R1 is phenyl,

phenyl substituted by R3 and/or R4,

naphthalenyl or

naphthalenyl substituted by R5 and/or R6,

R2 is phenyl,

phenyl substituted by R12 and/or R13,


naphthalenyl,

naphthalenyl substituted by R14 and/or R15,

aryl₂,

aryl₂ substituted by R16 and/or R17

or a radical selected from



R3 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy or benzyloxy,

- 88 -

- R4 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R5 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy or benzyloxy,
- R6 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,
- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

and the salts of these compounds with the proviso that the following compounds are excluded

- 4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(2-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-[6-(4-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,

- 89 -

N-{4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide,
 N-{4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide,
 N-{3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl}-acetamide,
 N-{4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide,
 4-Methyl-N-{4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl}-benzenesulfonamide,
 4-Methyl-N-{4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl}-benzenesulfonamide,
 N-{4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide,
 4-Methyl-N-{4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl}-benzenesulfonamide,
 N-{4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide, and
 N-{4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide.

3. Compounds of formula 1 according to claim 1 in which

- R1 is phenyl, 3-fluorophenyl, 4-fluorophenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-dimethoxyphenyl, 3-acetylphenyl, 4-acetylphenyl, 3-cyanophenyl, 4-phenoxyphenyl or naphthalen-1-yl,
 R2 is phenyl, phenyl substituted by R12 and/or R13, thiophenyl, naphthalenyl or 5-(1-methyl-5-trifluoromethyl-1H-pyrazol-3-yl)-thiophene-2-yl,
 R12 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine or 1-4C-alkoxycarbonyl,
 R13 is halogen, 1-4C-alkyl or 1-4C-alkoxy,

and the salts of these compounds with the proviso that the following compounds are excluded

4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide, and
 N-{4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl}-4-methyl-benzenesulfonamide.

4. Compounds of formula 1 according to claim 1 in which

- R1 is phenyl, 3-fluorophenyl, 4-fluorophenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-acetylphenyl or naphthalen-1-yl,
 R2 is 2-fluorophenyl, 3-fluorophenyl, 3,4-difluorophenyl, 2,4-difluorophenyl, 2,6-difluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluorophenyl, 3-chloro-4-methylphenyl, 3-bromophenyl, 3-methylphenyl, 4-methylphenyl, 4-isopropylphenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 4-cyanophenyl or 5-(1-methyl-5-trifluoromethyl-1H-pyrazol-3-yl)-thiophene-2-yl,

and the salts of these compounds with the proviso that the following compounds are excluded


4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide and
 4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

5. Compounds of formula 1 according to claim 1 in which

- R1 is phenyl substituted by R3 and/or R4 or naphthalenyl substituted by R5 and/or R6,

- 90 -

- R2 is phenyl,
 phenyl substituted by R12 and/or R13,
 naphthalenyl,
 naphthalenyl substituted by R14 and/or R15,
 aryl2,
 aryl2 substituted by R16 and/or R17
 or a radical selected from



- R3 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein
 R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and
 R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein
 R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,
 n is an integer from 0 to 4,
 m is an integer from 2 to 4,
 R4 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
 R5 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy,

- 91 -

1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R_{31})R_{32}$, $-CH_2CH(OH)CH_2N(R_{31})R_{32}$ or $-O-(CH_2)_m-N(R_{31})R_{32}$,

- R6 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,
- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

and the salts of these compounds.

6. Compounds of formula 1 according to claim 1 in which

- R1 is phenyl substituted in para or meta position by R3 or phenyl substituted in para and meta position by R3 and R4,
- R2 is phenyl substituted by R12 and/or R13,
- R3 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-

- 92 -

4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein

R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein

R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

n is an integer from 1 to 4,

m is an integer from 2 to 4,

R4 is fluorine, methyl or methoxy,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds.

7. Compounds of formula 1 according to claim 1 in which

R1 is phenyl substituted in para or meta position by R3,

R2 is phenyl substituted by R12 and/or R13,

R3 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein

R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein

R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

n is an integer from 1 to 4,

- 93 -

- m is an integer from 2 to 4,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- and the salts of these compounds.

8. Compounds of formula 1 according to claim 1 in which

- R1 is phenyl substituted in para position by R3 or phenyl substituted in para position by R3 and in meta position by R4,
- R2 is phenyl substituted by R12 and/or R13,
- R3 is morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylmethyl, 4-methylpiperazin-1-ylethyl, 4-methylpiperazin-1-ylpropyl, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 1-methylpiperidin-4-ylmethoxy, 1-methylpiperidin-4-ylmethyl, 1-methylpiperidin-4-ylethyl, 1-methylpiperidin-4-ylpropyl, 2-(1-methylpiperidin-4-yl)ethoxy, 3-(1-methylpiperidin-4-yl)propoxy, pyrrolidin-1-ylpropyl, pyrrolidin-1-ylethyl, pyrrolidin-1-ylmethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy, piperidin-1-ylmethyl, piperidin-1-ylethyl, piperidin-1-ylpropyl, 2-(piperidin-1-yl)ethoxy, 3-(piperidin-1-yl)propoxy,
- R4 is fluorine, methyl or methoxy,
- R12 is fluorine, chlorine, cyano, methyl, isopropyl, trifluoromethyl or methoxy,
- R13 is fluorine or chlorine,
- and the salts of these compounds.

9. Compounds of formula 1 according to claim 1 in which

- R1 is phenyl substituted in para or meta position by R3,
- R2 is phenyl substituted by R12 and/or R13,
- R3 is morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylmethyl, 4-methylpiperazin-1-ylethyl, 4-methylpiperazin-1-ylpropyl, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 1-methylpiperidin-4-ylmethoxy, 1-methylpiperidin-4-ylmethyl, 1-methylpiperidin-4-ylethyl, 1-methylpiperidin-4-ylpropyl, 2-(1-methylpiperidin-4-yl)ethoxy, 3-(1-methylpiperidin-4-yl)propoxy, pyrrolidin-1-ylpropyl, pyrrolidin-1-ylethyl, pyrrolidin-1-ylmethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy, piperidin-1-ylmethyl, piperidin-1-ylethyl, piperidin-1-ylpropyl, 2-(piperidin-1-yl)ethoxy, 3-(piperidin-1-yl)propoxy,
- R12 is fluorine, chlorine, cyano, methyl, isopropyl, trifluoromethyl or methoxy,

- 94 -

R13 is fluorine or chlorine,
and the salts of these compounds.

10. Compounds of formula 1 according to claim 1 in which

R1 is phenyl substituted in para position by R3 or phenyl substituted in para position by R3 and in meta position by R4,

R2 is 2-fluorophenyl, 4-fluorophenyl, 2,6-difluorophenyl, 2,4-difluorophenyl, 2-fluoro-4-methylphenyl, 2-fluoro-4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 4-methylphenyl, 4-methoxyphenyl or 3-chloro-4-fluorophenyl,

R3 is 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylethoxy, 4-methylpiperazin-1-ylpropoxy, morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 1-methylpiperidin-4-ylmethoxy, 2-(1-methylpiperidin-4-yl)ethoxy, 4-methylpiperazin-1-ylethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy or 3-pyrrolidin-1-ylpropyl,

R4 is fluorine,
and the salts of these compounds.

11. Compounds of formula 1 according to claim 1 in which

R1 is phenyl substituted in para or meta position by R3,

R2 is 2-fluorophenyl, 4-fluorophenyl, 2,6-difluorophenyl, 2,4-difluorophenyl, 4-methylphenyl, 4-methoxyphenyl or 3-chloro-4-fluorophenyl,

R3 is 2-morpholin-4-ylethoxy, 3-morpholin-4-ylpropoxy, 4-methylpiperazin-1-ylethoxy, 4-methylpiperazin-1-ylpropoxy, morpholin-4-ylmethyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, 1-methylpiperidin-4-ylmethoxy, 2-(1-methylpiperidin-4-yl)ethoxy, 4-methylpiperazin-1-ylethyl, 3-pyrrolidin-1-ylpropoxy, 2-pyrrolidin-1-ylethoxy or 3-pyrrolidin-1-ylpropyl,


and the salts of these compounds.

12. Compounds of formula 1 according to claim 1 in which

R1 is aryl1,
aryl1 substituted by R7 and/or R8,
R9 or
R10,


R2 is phenyl,
phenyl substituted by R12 and/or R13,
naphthalenyl,
naphthalenyl substituted by R14 and/or R15,
aryl2,
aryl2 substituted by R16 and/or R17
or a radical selected from

- 95 -



- Aryl1 furanyl, thiophenyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl or dibenzofuranyl,
- R7 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$, wherein
- R31 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and
- R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl or wherein
- R31 and R32 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,
- n is an integer from 0 to 4,
- m is an integer from 2 to 4,
- R8 is halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R9 is unsubstituted pyrrolyl, pyrazolyl, imidazolyl, indolyl, indazolyl, benzimidazolyl or benztriazolyl, or a radical selected from

- 96 -




wherein

R91 is hydroxyl, halogen, cyano, carboxyl, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, 1-4C-alkylcarbonyl, aminocarbonyl, mono- or di-1-4C-alkylaminocarbonyl, 1-4C-alkylcarbonylamino, phenoxy, benzyloxy, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkoxy, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkoxy, $-(CH_2)_n-N(R31)R32$, $-CH_2CH(OH)CH_2N(R31)R32$ or $-O-(CH_2)_m-N(R31)R32$,

R92 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R10 is a radical selected from



wherein

R101 is 1-4C-alkyl, 2,2,2-trifluoroethyl or 3,3,3-trifluoropropyl,

- 97 -

- R102 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,
- R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

and the salts of these compounds with the proviso that the following compounds are excluded

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide, and
 4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

13. Compounds of formula 1 according to claim 1 in which

- 98 -

- R1 is pyrid-3-yl, pyrid-4-yl, 2-methyl-2,3-dihydrobenzofuran-5-yl, benzo[1,3]dioxol-5-yl, 1-methyl-1H-pyrrol-3-yl, 4-methyl-thiophen-2-yl, 1-methyl-1H-pyrrol-2-yl, 1H-indol-5-yl, 1-methyl-1H-indol-3-yl, 1-methyl-1H-indol-5-yl, dibenzofuran-4-yl or 3,5-dimethyl-isoxazol-4-yl,
- R2 is phenyl substituted by R12 and/or R13 or naphthalenyl,
- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds with the proviso that the following compounds are excluded

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide, and
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide.

14. Compounds of formula 1 according to claim 1 in which

- R1 is 2-methyl-2,3-dihydrobenzofuran-5-yl, benzo[1,3]dioxol-5-yl, 1-methyl-1H-pyrrol-3-yl, 4-methyl-thiophen-2-yl, 1-methyl-1H-pyrrol-2-yl, 1H-indol-5-yl, 1-methyl-1H-indol-3-yl or 1-methyl-1H-indol-5-yl,
- R2 is phenyl substituted by R12 and/or R13 or naphthalenyl,
- R12 is fluorine, chlorine, cyano, methyl, trifluoromethyl or methoxy,
- R13 is fluorine, chlorine or methoxy,

and the salts of these compounds with the proviso that the compound

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide is excluded.

15. Compounds of formula 1 according to claim 1 in which

- R1 is 2-methyl-2,3-dihydrobenzofuran-5-yl, 1-methyl-1H-pyrrol-3-yl or 4-methylthiophen-2-yl, and
- R2 is 2,6-difluorophenyl,
- or
- R1 is 1H-indol-5-yl, and
- R2 is 2,6-difluorophenyl, 2,4-difluorophenyl, 2-fluoro-4-methylphenyl, 2-fluorophenyl or 3-fluorophenyl,
- or
- R1 is 1-methyl-1H-indol-5-yl or 1-methyl-1H-indol-3-yl, and
- R2 is 2,6-difluorophenyl, 2-fluorophenyl or 4-methoxyphenyl,
- or
- R1 is 1-methyl-1H-indol-5-yl, and
- R2 is 2-fluoro-4-methylphenyl,
- and the salts of these compounds.

- 99 -

16. Compounds of formula 1 according to claim 1 in which

either

R1 is 2-methyl-2,3-dihydrobenzofuran-5-yl, 1-methyl-1H-pyrrol-3-yl or 4-methylthiophen-2-yl, and

R2 is 2,6-difluorophenyl,

or

R1 is 1H-indol-5-yl, and

R2 is 2,6-difluorophenyl, 2,4-difluorophenyl, 2-fluorophenyl or 3-fluorophenyl,

or

R1 is 1-methyl-1H-indol-5-yl or 1-methyl-1H-indol-3-yl, and

R2 is 2,6-difluorophenyl, 2-fluorophenyl or 4-methoxyphenyl,

and the salts of these compounds.

17. Compounds of formula 1 according to claim 1 in which

R1 is R11,

R2 is phenyl,

phenyl substituted by R12 and/or R13,


naphthalenyl,

naphthalenyl substituted by R14 and/or R15,

aryl²,


aryl² substituted by R16 and/or R17

or a radical selected from



R11 is a radical selected from

- 100 -



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R112)R113$ or $-CH_2CH(OH)CH_2N(R112)R113$, wherein

R112 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R113 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R112 and R113 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

R14 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R15 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

Aryl2 furanyl, thiophenyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, benzofuranyl, benzothiophenyl, 2,3-dihydrobenzofuranyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzo[1,3]dioxolyl, 2,3-dihydrobenzo[1,4]-

- 101 -

dioxinyl, quinazolinyl, quinoxalinyl, cinnolinyl, quinolinyl, isoquinolinyl, phthalazinyl, indanyl, indolyl or indazolyl,

R16 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R17 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,


and the salts of these compounds.

18. Compounds of formula 1 according to claim 1 in which

R1 is R11,

R2 is phenyl substituted by R12 and/or R13,

R11 is a radical selected from



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R112)R113$ or $-CH_2CH(OH)CH_2N(R112)R113$, wherein

R112 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R113 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R112 and R113 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

- 102 -


- R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,
- R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,
- and the salts of these compounds.

19. Compounds of formula 1 according to claim 1 in which

R1 is R11,

R2 is phenyl substituted by R12 and/or R13,

R11 is a radical selected from



wherein

R111 is 1-(1-4C-alkyl)-pyrrolidin-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-pyrrolidin-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-piperid-4-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-2-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-3-yl-1-4C-alkyl, 1-(1-4C-alkyl)-azepan-4-yl-1-4C-alkyl, $-(CH_2)_p-N(R112)R113$ or $-CH_2CH(OH)CH_2N(R112)R113$, wherein

R112 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, and

R113 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-2-4C-alkyl, or wherein

R112 and R113 together and with inclusion of the nitrogen atom to which they are bonded form a pyrrolidin-, piperidin-, 4-hydroxy-piperidin-, piperazin-, 4-(1-4C-alkyl)piperazin-, [1,4]diazepan-, 4-(1-4C-alkyl)-[1,4]diazepan-, morpholin-, thiomorpholin- or an azepan-ring,

p is an integer from 1 to 4,

R114 is hydrogen, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy completely or predominantly substituted by fluorine, amino or mono- or di-1-4C-alkylamino,

R12 is hydroxyl, halogen, cyano, nitro, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkylamino,

R13 is hydroxyl, halogen, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino or mono- or di-1-4C-alkyl-amino,

and the salts of these compounds.


- 103 -

20. Compounds of formula 1 according to claim 1 in which

R1 is R11,

R2 is phenyl substituted by R12 and/or R13,

R11 is a radical selected from



wherein

R111 is pyrrolidin-1-ylethyl, pyrrolidin-1-ylpropyl, piperidin-1-ylethyl, piperidin-1-ylpropyl, azepan-1-ylethyl, azepan-1-ylpropyl, (4-methyl-piperazin-1-yl)ethyl, (4-methyl-piperazin-1-yl)propyl, morpholin-4-ylethyl, morpholin-4-ylpropyl, (1-methyl-piperidin-4-yl)propyl, (1-methyl-piperidin-4-yl)ethyl, (1-methyl-piperidin-4-yl)methyl, dimethylaminoethyl, dimethylaminopropyl, dimethylaminobutyl, diethylaminoethyl, diethylaminopropyl or diethylaminobutyl,

R114 is hydrogen or fluorine,

R12 is fluorine, chlorine, cyano, methyl, isopropyl, trifluoromethyl or methoxy,

R13 is fluorine or chlorine,


and the salts of these compounds.

21. Compounds of formula 1 according to claim 1 in which

R1 is R11,

R2 is 2-fluorophenyl, 4-fluorophenyl, 2,6-difluorophenyl, 2,4-difluorophenyl, 4-methylphenyl, 4-methoxyphenyl, 3-fluoro-4-methoxyphenyl, 2-fluoro-4-methylphenyl, or 3-chloro-4-fluorophenyl,

R11 represents the following radical



wherein

R111 is pyrrolidin-1-ylethyl, pyrrolidin-1-ylpropyl, (4-methyl-piperazin-1-yl)ethyl, (4-methyl-piperazin-1-yl)propyl, (1-methyl-piperidin-4-yl)ethyl, (1-methyl-piperidin-4-yl)methyl, dimethylaminoethyl, dimethylaminopropyl, diethylaminopropyl or diethylaminobutyl,

and the salts of these compounds.

22. Compounds according to claim 1 for use in the treatment of diseases.

- 104 -

23. Pharmaceutical compositions containing one or more compounds according to claim 1 together with the usual pharmaceutical auxiliaries and/or excipients.

24. Use of compounds according to claim 1 for the production of pharmaceutical compositions for the treatment of acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock. The compounds according to the invention are also useful in the treatment and/or prevention of T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrheic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

25. A method for treating acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock, T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrheic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis in a patient comprising administering to said patient a therapeutically effective amount of a compound according to claim 1.

26. A compound selected from
4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

- 105 -

N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-[6-(2-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-(3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl)-acetamide,
N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
or a salt thereof for use in the treatment of diseases.

27. Pharmaceutical composition containing one or more compounds selected from
4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-[6-(2-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-(3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl)-acetamide,
N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,

- 106 -

4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
or a salt thereof together with the usual pharmaceutical auxiliaries and/or excipients.

28. Use of a compound selected from

4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-[6-(2-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Benzyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-(3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl)-acetamide,
N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,

- 107 -

4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 or a salt thereof for the production of pharmaceutical compositions for the treatment of acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock. The compounds according to the invention are also useful in the treatment and/or prevention of T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

29. A method for treating acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock, T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis in a patient comprising administering to said patient a therapeutically effective amount of a compound selected from 4-Methyl-N-[4-(6-naphthalen-2-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
 N-[4-[6-(Bis-trifluoromethyl-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
 4-Methyl-N-[4-(6-phenyl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

- 108 -

4-Methyl-N-[4-(6-naphthalen-1-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-[6-(2-Benzoyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Benzoyloxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3,4-Dimethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(4-Hydroxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-(3-[6-[4-(Toluene-4-sulfonylamino)-phenylamino]-pyrimidin-4-yl]-phenyl)-acetamide,
N-[4-[6-(2-Methoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(3-nitro-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-trifluoromethoxy-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(4-Cyano-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-[6-(4-morpholin-4-yl-phenyl)-pyrimidin-4-ylamino]-phenyl]-benzenesulfonamide,
N-[4-[6-(3-Amino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-[6-(3-Dimethylamino-phenyl)-pyrimidin-4-ylamino]-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide
or a salt thereof.

30. A compound selected from

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide, and
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
or a salt thereof for use in the treatment of diseases.

31. Pharmaceutical compositions containing one or more compounds selected from N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,

- 109 -

4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
or a salt thereof together with the usual pharmaceutical auxiliaries and/or excipients.

32. Use of compounds selected from

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,
4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,
or a salt thereof for the production of pharmaceutical compositions for the treatment of acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock. The compounds according to the invention are also useful in the treatment and/or prevention of T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

33. A method for treating acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure,

- 110 -

chronic obstructive pulmonary disease, CNS diseases such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious diseases such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, or traumatic shock, T-cell mediated acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I or II and the disorders associated therewith, respiratory diseases such as asthma or inflammatory lung injury, inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis in a patient comprising administering to said patient a therapeutically effective amount of a compound selected from

N-[4-(6-Benzo[1,3]dioxol-5-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,

4-Methyl-N-[4-(6-pyridin-4-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

N-[4-(2',4'-Dimethoxy-[4,5']bipyrimidinyl-6-ylamino)-phenyl]-4-methyl-benzenesulfonamide,

4-Methyl-N-[4-(6-pyridin-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

N-[4-(6-Benzofuran-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,

4-Methyl-N-[4-(6-thiophen-3-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

N-[4-(6-Dibenzofuran-4-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide,

N-[4-(6-Benzo[b]thiophen-2-yl-pyrimidin-4-ylamino)-phenyl]-4-methyl-benzenesulfonamide, and

4-Methyl-N-[4-(6-quinolin-8-yl-pyrimidin-4-ylamino)-phenyl]-benzenesulfonamide,

or a salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2005/050206

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/42 C07D409/14 C07D403/04 C07D405/04 A61K31/505
A61K31/506 A61P35/00 A61P37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 2005/026129 A1 (AXXIMA PHARMACEUTICALS AG, GERMANY) 24 March 2005 (2005-03-24) see e.g. compounds 1, 2, 5 (p. 124), 23 (p. 125), 34 (p. 126) and 155 (p. 132) of table 2	1-33
A	US 5 821 246 A (BROWN ET AL) 13 October 1998 (1998-10-13) see abstract, claim 1, example 12 and compound 26 from example 15, table I	1-33
A	WO 99/24440 A (PFIZER PRODUCTS INC; MUNCHHOF, MICHAEL, JOHN; SOBLOV-JAYNES, SUSAN, B) 20 May 1999 (1999-05-20) see claim 1, 1st compound on p. 5 and abstract	1-33

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 June 2005

Date of mailing of the international search report

16/06/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Traegler-Goeldel, M

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2005/050206

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 25, 28, 29, 32 and 33 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2005/050206

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2005026129	A1	24-03-2005	NONE	
US 5821246	A	13-10-1998	AT 175962 T	15-02-1999
			AU 703328 B2	25-03-1999
			AU 3813095 A	06-06-1996
			CA 2200871 A1	23-05-1996
			DE 69507505 D1	04-03-1999
			DE 69507505 T2	02-06-1999
			DK 790986 T3	13-09-1999
			EP 0790986 A1	27-08-1997
			ES 2128092 T3	01-05-1999
			FI 971970 A	07-05-1997
			WO 9615118 A1	23-05-1996
			IL 115959 A	20-06-2004
			JP 10508616 T	25-08-1998
			NO 972152 A	12-05-1997
			NZ 294917 A	25-02-1999
			ZA 9509572 A	13-05-1996
WO 9924440	A	20-05-1999	AP 976 A	12-06-2001
			AU 9454198 A	31-05-1999
			BG 104412 A	28-02-2001
			BR 9814018 A	26-09-2000
			CA 2309690 A1	20-05-1999
			CN 1280580 A	17-01-2001
			CZ 20001709 A3	12-12-2001
			EP 1028964 A1	23-08-2000
			HR 20000286 A1	31-12-2000
			HU 0100287 A2	28-09-2001
			WO 9924440 A1	20-05-1999
			ID 23978 A	14-06-2000
			JP 2001522853 T	20-11-2001
			MA 24694 A1	01-07-1999
			NO 20002162 A	10-07-2000
			NZ 520093 A	26-03-2004
			OA 11376 A	28-01-2004
			PL 340589 A1	12-02-2001
			SK 6652000 A3	09-05-2002
			TW 593321 B	21-06-2004
			US 6492383 B1	10-12-2002
			ZA 9810253 A	10-05-2000